

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
1 November 2001 (01.11.2001)

PCT

(10) International Publication Number  
**WO 01/81334 A2**

(51) International Patent Classification<sup>7</sup>: **C07D 401/00**

(21) International Application Number: **PCT/JP01/03378**

(22) International Filing Date: **20 April 2001 (20.04.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:  
**2000-120976** **21 April 2000 (21.04.2000)** **JP**

(71) Applicant (for all designated States except US): **SUN-  
TORY LIMITED** [JP/JP]; 1-40, Dojimahama 2-chome,  
Kita-ku, Osaka-shi, Osaka 530-8203 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **IMOTO, Masahiro**  
[JP/JP]; 15-501, Naka-machi 3-chome, Tatebayashi-shi,  
Gunma 374-0029 (JP). **IWANAMI, Tatsuya** [JP/JP];  
16-16, Dai-cho, Ashikaga-shi, Tochigi 326-0056 (JP).

**AKABANE, Minako** [JP/JP]; 1-37-201, Sojiji 1-chome,  
Ibaraki-shi, Osaka 567-0801 (JP). **TANI, Yoshihiro**  
[JP/JP]; 9-38, Hoshimi-cho, Ibaraki-shi, Osaka 567-0843  
(JP).

(74) Agent: **KUSAMA, Osamu**; KUSAMA PATENT  
OFFICE, 7F Iwata Bldg., 5-12, Iidabashi 4-chome, Chiy-  
oda-ku, Tokyo 102-0072 (JP).

(81) Designated States (national): **AU, CA, CN, KR, US.**

(84) Designated States (regional): European patent (AT, BE,  
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,  
NL, PT, SE, TR).

**Published:**


— without international search report and to be republished  
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.



**WO 01/81334 A2**

(54) Title: **CYCLIC AMIDINE COMPOUNDS**



(57) Abstract: There is provided cyclic amidine compounds of the following formula (I) wherein: A<sup>1</sup> and A<sup>2</sup> are hydrogen atom, optionally substituted alkyl group; optionally substituted aryl group; or optionally substituted heterocyclic group; and X is -C(R<sup>1</sup>,R<sup>2</sup>)-C(R<sup>3</sup>,R<sup>4</sup>)-, -C(R<sup>5</sup>)=C(R<sup>6</sup>)-, -C(R<sup>7</sup>,R<sup>8</sup>)-C(R<sup>9</sup>,R<sup>10</sup>)-C(R<sup>11</sup>,R<sup>12</sup>)-, or -C(R<sup>13</sup>,R<sup>14</sup>)-C(R<sup>15</sup>,R<sup>16</sup>)-NH- (wherein, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup> and R<sup>16</sup> are hydrogen atom; halogen atom; optionally substituted alkyl group; optionally substituted aryl group; or optionally substituted heterocyclic group; or pharmaceutically acceptable salts thereof. These compounds have good

affinity for  $\alpha 4 \beta 2$  nicotinic acetylcholine receptors and activate the same to thereby exert a preventive or therapeutic effect on cerebral dysfunction.

## DESCRIPTION

## CYCLIC AMIDINE COMPOUNDS

## 5 TECHNICAL FIELD

The present invention relates to compounds showing affinity for nicotinic acetylcholine receptors and activating the same. The compounds of the present invention are useful for preventing or treating of neurodegenerative diseases such as  
10 Alzheimer's disease and Parkinson's disease, dementia such as cerebrovascular dementia, motor ataxia such as Tourette's syndrome, neurosis during chronic cerebral infarction stage, neuropathy and mental disorder such as anxiety and schizophrenia and cerebral dysfunction caused by cerebral injury.

15

## BACKGROUND ART

It has been widely known that nicotine exerts a wide variety of pharmacological effects. These include, for example, cholinergic nervous activation as the effect on central nervous  
20 systems such as facilitation of acetylcholine release [De sarno P. & Giacobini E., *J. Neurosci. Res.*, 22, 194-200 (1984)], and further, activation effect on monoaminergic nervous systems [Levin E. D. & Simon B. B., *Psychopharmacology*, 138, 217-230 (1998)].

25

It has been also reported that nicotine possesses lots of very useful cerebral function improving effects such as increasing cerebral blood flow and glucose uptake rate in brain [Decker M. W. et al., *Life Sci.*, 56, 545-570 (1995)].

30

It has been further reported that nicotine inhibits amyloid formation of  $\beta$ -peptides which is believed to be the cause of neuronal cell death during Alzheimer's disease [Salomon A. R. et al., *Biochemistry*, 35, 13568-13578 (1996)], and have cell

protective effects on neuronal cell death induced by  $\beta$ -amyloid (A $\beta$ ) [Kihara T. et al., *Ann. Neurol.*, 42, 156-163 (1997)]. Recent studies suggest the possibility of nicotine being a remedy for the inflammatory colitis [Sandborn W. J. et al., *Ann. Intern. Med.*, 126, 364 (1997)].

On the other hand, it is acknowledged that in the patients of Alzheimer's disease, the degeneration of acetylcholinergic neurons known to be one of the important nervous systems responsible for cognition such as attention, learning, memory and recognition, is altered and thus nicotinic acetylcholine receptors in the cerebral cortex and hippocampus are drastically decreased [Nordberg A. et al., *J. Neurosci. Res.*, 31, 103-111 (1992)].

It is reported the possibility of the useful treatment for Alzheimer's disease by activating nicotinic acetylcholine receptors to be recovered the acetylcholine nervous systems mechanism by agonists or modulators of nicotinic acetylcholine receptors [Newhouse P. A. et al., *Psychopharmacology*, 95, 171-175 (1988)].

The nicotinic acetylcholine receptors belong to the ion channel neurotransmitter receptors composed of five subunits. That is, agonists such as acetylcholine, nicotine and the like are bound to receptors to activate and open the channels thereof, thus causing the influx of cationic ion such as sodium ion from extracellular to result the cell excitation [Galzi J. L. & Changeux J. P., *Neuropharmacology*, 34, 563-582 (1995)]. The aforementioned agonists such as acetylcholine, nicotine and the like show its effect by binding to the specific site existing in  $\alpha$  subunit so-called agonist binding site.

It is known, on the other hand, that compounds such as galantamine and so on which activate cells by potentiating the effects of acetylcholine, have no agonist effect at nicotinic

acetylcholine receptors directly. These compounds show their effects through allosteric site which is clearly different from the agonist binding sites [Schrattenholz A. et al., *Mol. Pharmacol.*, 49, 1-6 (1996)].

5           Mentioned above, compounds capable to activate nicotinic acetylcholine receptors indirectly are called modulators and it is expected to be the practical medicines for treatment of the various neurological diseases [Lin N. -H & Meyer M. D., *Exp. Opin. Thr. Patents*, 8, 991-1015 (1998)].

10           The terms "agonists" and "modulators" are used in these definitions in the present specification.

          The nicotinic acetylcholine receptors are believed to participate not only in Alzheimer's disease, but also in neurodegenerative diseases such as Parkinson's disease, and many  
15 of the neuroses and psychoses such as dementia, anxiety, schizophrenia and so on [Barrantes F. J., in *The Nicotinic Acetylcholine Receptor*, ed. Barrantes F. J., Springer, 1997, p175-212; Lena C. & Changeux J. -P., *J. Physiol. (Paris)*, 92, 63-74 (1998)].

20           Especially, since it is known that cerebral blood flow of the patients suffering from cerebrovascular dementia caused by cerebral infarction is decreased [Takagi Shigeharu, *Gendai Iryo*, 28, 1157-1160 (1996); Tachibana H. et al., *J. Gerontol.*, 39, 415-423 (1984)], there seems to be the possibility of agonists of  
25 nicotinic acetylcholine receptors or the modulators possessing cerebral blood flow increasing effect to be applied to the medicines in this area of treatment. Furthermore, recent study revealed that agonists of nicotinic acetylcholine receptors and the modulators thereof show analgesic activities [Bannon A. W. et  
30 al., *Science*, 279, 77-81 (1998)].

          Nicotine itself surely affects as the agonist of nicotinic acetylcholine receptors. For example, after administration of

nicotine to the patients of Alzheimer's disease, the recoveries of their attention or the short-term memory were observed, and also the symptoms of their disease were improved [Newhouse P. A. et al., *Drugs & Aging*, 11, 206-228 (1997)]. Nevertheless, nicotine also possesses disadvantages such as widely recognized addiction, as well as low bioavailability and severe side effects to the cardiovascular systems.

Therefore, there have been great expectation to develop nicotinic acetylcholine receptors agonists or modulators as medicines in place of nicotine which has no addiction, high bioavailability, and less side effects on cardiovascular systems [Maelicke A. & Albuquerque E. X., *Drug Discovery Today*, 1, 53-59 (1996); Holladay M. W. et al., *J. Med. Chem.*, 40, 4169-4194 (1997)].

There are some subtypes known as the nicotinic acetylcholine receptors [Shacka J. J. & Robinson S. E. T., *Med. Chem. Res.*, 1996, 444-464], and mainly  $\alpha 4 \beta 2$  subtype receptors exist in central nervous systems. Furthermore, there exist  $\alpha 1 \beta 1 \gamma \delta$  (or  $\alpha 1 \beta 1 \epsilon \delta$ ) subtype receptors in the neuromuscular junction of motor neurons, and  $\alpha 3 \beta 4$  subtype receptors in ganglion of autonomic nervous systems and adrenal.


The activation of the cholinergic nervous systems and increasing effect of cerebral blood flow are believed to occur though  $\alpha 4 \beta 2$  subtype receptors in central nervous systems, and above mentioned effects of nicotine on cardiovascular system are induced by affecting receptor subtypes exist in peripheral nervous system.

Therefore, it may be extremely useful as medicines having no side effects to develop compounds which have no affinity at  $\alpha 1 \beta 1 \gamma \delta$  subtype nor  $\alpha 3 \beta 4$  subtype receptors, but selectively affects  $\alpha 4 \beta 2$  subtype receptors.

In these circumstances, there have been many proposals to

develop selective agonists or modulators at nicotinic acetylcholine receptors of central nervous system as practical medicines. These include, for example, the compound such as ABT-418 [Arneric S. P. et al., *J. Pharmacol. Exp. Ther.*, 270, 310-318 (1994); Decker M. W. et al., *J. Pharmacol. Exp. Ther.*, 270, 319-328 (1994)], ABT-089 [Sullivan J. P. et al., *J. Pharmacol. Exp. Ther.*, 283, 235-246 (1997); Decker M. W. et al., *J. Pharmacol. Exp. Ther.*, 283, 247-258 (1997)], GTS-21 [Arendash G. W. et al., *Brain Res.*, 674, 252-259 (1995); Briggs C. A. et al., *Pharmacol. Biochem. Behav.*, 57, 231-241 (1997)], RJR-2403 [Bencherif M. et al., *J. Pharmacol. Exp. Ther.*, 279, 1413-1421 (1996); Lippiello P. M. et al., *J. Pharmacol. Exp. Ther.*, 279, 1422-1429 (1996)], SIB-1508Y [Cosford N. D. P. et al., *J. Med. Chem.*, 39, 3235-3237 (1996); Lloyd. G. K. et al., *Life Sci.*, 62, 1601-1606 (1995)], SIB-1553A [Lloyd. G. K. et al., *Life Sci.*, 62, 1601-1606 (1995)] and so on.

In European Patent Publication EP679397-A2, substituted amine derivatives represented by the following formula were proposed for the medicines for prevention and treatment of cerebral dysfunction.



in which,

R represents hydrogen, optionally substituted acyl, alkyl, aryl, aralkyl, heteroaryl or heteroarylalkyl radicals;

A represents a monofunctional group of the hydrogen, acyl, alkyl or aryl series or represents a bifunctional group which is linked to the radical Z;

E represents an electron-withdrawing radical;

X represents the -CH= or =N- radicals, it being possible for the -CH= radical to be linked to the Z radical

instead of an H atom;


Z represents a monofunctional group of the alkyl, -O-R, -S-R or -NR<sub>2</sub> series or represents a bifunctional group which is linked to the A radical or the X radical.

5

However, the structure of the compounds disclosed in said patent publication is clearly different from that of the compounds disclosed by the present patent application, and there is no description in the above-mentioned patent publication that  
10 these compounds can selectively activate  $\alpha 4\beta 2$  nicotinic acetylcholine receptors.

On the other hand, it is confirmed that "imidacloprid", as a pesticide, electrophysiologically affects as partial agonist at nicotinic acetylcholine receptors of PC12 cell [Nagata K. et al.,  
15 *J. Pharmacol. Exp. Ther.*, 285, 731-738 (1998)], and imidacloprid itself or its metabolites and their analogues possess affinity to the nicotinic acetylcholine receptors in mouse brain [Lee Chao S. & Casida E., *Pestic. Biochem. Physiol.*, 58, 77-88 (1997); Tomizawa T. & Casida J. E., *J. Pharmacol.*, 127, 115-122 (1999);  
20 Latli B. et al., *J. Med. Chem.*, 42, 2227-2234 (1999)], however, there is no report of the imidacloprid derivatives selectively activating  $\alpha 4\beta 2$  nicotinic acetylcholine receptors. Furthermore, the structure of the imidacloprid itself or its metabolites and their analogues is clearly different from that of the compounds  
25 disclosed by the present patent application.

Japanese Laid-open Patent Publication Number Hei 10-226684 disclosed [N-(pyridinylmethyl)heterocyclic]ylideneamine compounds represented by the following formula, pharmaceutically acceptable  
30 salts and prodrugs thereof.




in which,

A represents the -CH(R)-;

R<sup>3</sup> represents a hydrogen atom or an optionally substituted  
 5 C<sub>1</sub>-C<sub>6</sub> alkyl; and

B represents the group of the following formula:



It is disclosed that these compounds possess weak affinity  
 to nicotinic receptors; however, there is no description that  
 10 these compounds have selective activating effect at  $\alpha 4\beta 2$   
 nicotinic acetylcholine receptors of central nervous systems and  
 act as agonists or modulators of nicotinic acetylcholine  
 receptors. Furthermore, the structure of these compounds is  
 clearly different from that of the compounds disclosed by the  
 15 present invention.

As mentioned above, there had been many attempts to  
 develop agonists or modulators selectively activating  $\alpha 4\beta 2$   
 nicotinic acetylcholine receptors of central nervous systems via  
 oral administration, but none were satisfactory.

20

#### DISCLOSURE OF THE INVENTION

Therefore, the present invention provides therapeutic or  
 preventing agents for treatment of diseases which may be  
 prevented or cured by activating nicotinic acetylcholine  
 25 receptors, having the capabilities of binding selectively with  
 $\alpha 4\beta 2$  nicotinic acetylcholine receptor of central nervous systems,




and having no undesirable side effects in cardiovascular systems such as hypertension or tachycardia.

More specifically, the present invention provides medicaments for preventing or treating various diseases, which may be prevented or cured by activating nicotinic acetylcholine receptors, such as dementia, senile dementia, presenile dementia, Alzheimer's disease, Parkinson's disease, cerebrovascular dementia, AIDS-related dementia, dementia in Down's syndrome, Tourette's syndrome, neurosis during chronic cerebral infarction stage, cerebral dysfunction caused by cerebral injury, anxiety, schizophrenia, depression, Huntington's disease, pain and so on.

Through extensive investigations of researching compounds having the capabilities of binding selectively with  $\alpha 4\beta 2$  nicotinic acetylcholine receptors of central nervous systems, the present inventors discovered that the compounds represented by the formula (I) mentioned below and pharmaceutically acceptable salts thereof possess high affinity for nicotinic acetylcholine receptors in central nervous systems, and activate said receptors as agonists or modulators.

Accordingly, as one aspect of the present invention, it is provided cyclic amidine compounds represented by the following formula (I):



wherein:

$A^1$  and  $A^2$  are hydrogen atom, optionally substituted alkyl group; optionally substituted aryl group; or optionally substituted heterocyclic group; and

X is  $-C(R^1, R^2)-C(R^3, R^4)-$ ,  $-C(R^5)=C(R^6)-$ ,  $-C(R^7, R^8)-C(R^9, R^{10})-$

C(R<sup>11</sup>,R<sup>12</sup>)-, or -C(R<sup>13</sup>,R<sup>14</sup>)-C(R<sup>15</sup>,R<sup>16</sup>)-NH- (wherein, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup> and R<sup>16</sup> are hydrogen atom; halogen atom; optionally substituted alkyl group; optionally substituted aryl group; or optionally substituted  
5 heterocyclic group;  
or pharmaceutically acceptable salts thereof.

Still another aspect of the present invention, it is provided activator agents for  $\alpha 4\beta 2$  nicotinic acetylcholine  
10 receptors containing cyclic amidine compounds of the formula (I) or pharmaceutically acceptable salt thereof as active ingredients.

As still further aspect of the present invention, it is provided that the use of cyclic amidine compounds of the formula (I) or pharmaceutically acceptable salt thereof for treating or  
15 preventing of cerebral circulation disease, neurodegenerative disease and the like.

#### BEST MODE FOR CARRYING OUT THE INVENTION

Examples of the pharmaceutically acceptable salt include  
20 an inorganic acid salt such as hydrochloric acid salt, hydrobromic acid salt, sulfuric acid salt, phosphoric acid salt and the like, and an organic acid salt such as fumaric acid salt, maleic acid salt, oxalic acid salt, citric acid salt, tartaric acid salt, malic acid salt, lactic acid salt, succinic acid salt,  
25 benzoic acid salt, methanesulfonic acid salt, p-toluenesulfonic acid salt and the like.

The groups represented by "A<sup>1</sup>" and "A<sup>2</sup>" in the compound of formula (I) are hydrogen atom, optionally substituted alkyl group, optionally substituted aryl group or optionally substituted  
30 heterocyclic group, and preferable examples of said optionally substituted alkyl group include methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, tert-butyl and the like.

Suitable substituent of substituted alkyl group may include optionally substituted aryl group or optionally substituted heterocyclic group, and therefore, examples of said substituted alkyl group include benzyl, (2-pyridyl)methyl, (3-pyridyl)methyl, (2-chloro-3-pyridyl)methyl, (6-chloro-3-pyridyl)-methyl, (6-fluoro-3-pyridyl)methyl, (5-bromo-3-pyridyl)methyl, (2,6-dichloro-3-pyridyl)methyl, (5,6-dichloro-3-pyridyl)methyl, (2,6-dichloro-3-pyridyl)methyl, (6-methyl-3-pyridyl)methyl, (6-ethoxy-3-pyridyl)methyl, (5-pyrimidyl)methyl, (3-quinolyl)-methyl, (3-furanyl)methyl, (tetrahydro-3-furanyl)-methyl, (3-thienyl)-methyl, (3,5-dimethylisoxazolyl)methyl, 1-(6-chloro-3-pyridyl)-ethyl, 2-(6-chloro-3-pyridyl)ethyl and the like.


The preferable examples of aryl group of said optionally substituted aryl group represented by "A<sup>1</sup>" and "A<sup>2</sup>" may include phenyl, naphthyl and the like. Suitable substituent of substituted aryl group may include C<sub>1</sub>-C<sub>4</sub> lower alkyl group, hydroxyl group, amino group, halogen atom and the like, and therefore, examples of said substituted aryl group include methylphenyl, hydroxyphenyl, aminophenyl, chlorophenyl, dichlorophenyl and the like.

The term "heterocyclic group" represented by "A<sup>1</sup>" and "A<sup>2</sup>" may be 5 or 6 membered heterocyclic group or condensed heterocyclic group thereof containing the same or different 1 to 3 hetero atom(s) such as sulfur, nitrogen, oxygen atom(s), and examples include thiophene, furan, pyran, pyrrole, pyrazole, pyridine, pyrimidine, pyrazine, pyridazine, imidazole, oxazole, isoxazole, thiazole, isothiazole, quinoline, isoquinoline, indole, azaindole, tetrahydropyrimidine and the like.

Suitable substituent of substituted heterocyclic group may include C<sub>1</sub>-C<sub>4</sub> lower alkyl, halogen atom and the like, and therefore, examples of said substituted heterocyclic group may be 2-methylpyridine, 6-methylpyridine, 2-chloropyridine, 2-

fluoropyridine, 2-bromopyridine, 3-bromopyridine, 2,3-dichloropyridine, 2-chloropyrimidine, 2-chlorothiazole, 3,5-dimethylisoxazole and the like.

The group represented by "X" is the partial component of the bond as following;



wherein,  $\text{R}^1$  to  $\text{R}^{16}$  are hydrogen atom; halogen atom; optionally substituted alkyl group; optionally substituted aryl group; or optionally substituted heterocyclic group.

The term "halogen atom" represented by  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^3$ ,  $\text{R}^4$ ,  $\text{R}^5$ ,  $\text{R}^6$ ,  $\text{R}^7$ ,  $\text{R}^8$ ,  $\text{R}^9$ ,  $\text{R}^{10}$ ,  $\text{R}^{11}$ ,  $\text{R}^{12}$ ,  $\text{R}^{13}$ ,  $\text{R}^{14}$ ,  $\text{R}^{15}$  and  $\text{R}^{16}$  may include fluorine, chlorine, bromine and iodine.

The term "optionally substituted alkyl group" may include methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, tert-butyl and the like.

Suitable substituent of substituted alkyl group may include optionally substituted aryl group or optionally substituted heterocyclic group, and therefore, examples of said substituted alkyl group include benzyl, (2-pyridyl)methyl, (3-pyridyl)methyl, (2-chloro-3-pyridyl)methyl, (6-chloro-3-pyridyl)-methyl, (6-fluoro-3-pyridyl)methyl, (5-bromo-3-pyridyl)methyl, (2,6-dichloro-3-pyridyl)methyl, (5,6-dichloro-3-pyridyl)methyl, (2,6-dichloro-3-pyridyl)methyl, (6-methyl-3-pyridyl)methyl, (6-ethoxy-3-pyridyl)methyl, (5-pyrimidyl)methyl, (3-quinolyl)methyl, (3-furanyl)methyl, (tetrahydro-3-furanyl)methyl, (3-thienyl)-methyl, (3,5-dimethylisoxazolyl)methyl, 1-(6-chloro-3-pyridyl)-ethyl, 2-(6-chloro-3-pyridyl)ethyl and the like.

The term "optionally substituted aryl group" for the groups  $\text{R}^1$  to  $\text{R}^{16}$  may be non-substituted phenyl group or phenyl

group which is substituted by halogen atom, or C<sub>1</sub>-C<sub>4</sub> lower alkyl such as methyl, ethyl and the like, and therefore, examples of substituted phenyl group may include methylphenyl, chlorophenyl, dichlorophenyl and the like.

5           The term "heterocyclic group" for the groups R<sup>1</sup> to R<sup>16</sup> may be 5 or 6 membered heterocyclic group containing the same or different 1 to 3 hetero atom(s) such as sulfur, nitrogen, oxygen atom(s), and examples include thiophene, furan, pyran, pyrrole, pyrazole, pyridine, pyrimidine, pyrazine, pyridazine, imidazole,  
10   oxazole, isoxazole, thiazole, isothiazole, quinoline, isoquinoline, tetrahydropyrimidine and the like.

Suitable substituent of substituted heterocyclic group may include C<sub>1</sub>-C<sub>4</sub> lower alkyl, halogen atom and the like, and therefore, examples of said substituted heterocyclic group may be  
15   2-methylpyridine,      3-methylpyridine,      2-chloropyridine,      2-fluoropyridine,      2-bromopyridine,      3-bromopyridine,      2,3-dichloropyridine,      4-chloropyrimidine,      2-chlorothiazole,      3-methylisoxazole and the like.

20           The following are examples of cyclic amidine compounds of the formula (I).

- Compound 1: 2-(6-chloro-3-pyridyl)-2-imidazoline;  
Compound 2: 2-(6-chloro-3-pyridyl)-1,4,5,6-tetrahydropyrimidine;  
Compound 3: 2-(6-chloro-3-pyridyl)-1-methyl-2-imidazoline;  
25   Compound 4: 2-(6-chloro-3-pyridyl)-1-methyl-1,4,5,6-tetrahydropyrimidine;  
Compound 5: 1-(6-chloro-3-pyridyl)methylimidazole;  
Compound 6: 2-(6-chloro-3-pyridyl)imidazole;  
Compound 7: 2-(6-chloro-3-pyridyl)methyl-2-imidazoline;  
30   Compound 8: 2-(6-chloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;  
Compound 9: 2-(6-chloro-3-pyridyl)methyl-1-methyl-2-imidazoline;

- Compound 10: 2-(6-chloro-3-pyridyl)methyl-1-methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 11: 1-(6-chloro-3-pyridyl)methyl-2-methyl-2-imidazoline;
- Compound 12: 1-(6-chloro-3-pyridyl)methyl-4,4-dimethyl-2-imidazoline;
- Compound 13: 2-(tetrahydrofuran-3-yl)-1,4,5,6-tetrahydropyrimidine;
- Compound 14: 2-(tetrahydrofuran-3-yl)-2-imidazoline;
- Compound 15: 2-(tetrahydrofuran-3-yl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 16: 2-(5-bromo-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 17: 2-(5-bromo-3-pyridyl)methyl-2-imidazoline;
- Compound 18: 2-(3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 19: 2-(3-pyridyl)methyl-2-imidazoline;
- Compound 20: 2-(3-aminophenyl)-1,4,5,6-tetrahydropyrimidine;
- Compound 21: 2-(3-quinolyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 22: 2-(2-chloro-5-thiazolyl)-1,4,5,6-tetrahydropyrimidine;
- Compound 23: 2-(3-quinolyl)methyl-2-imidazoline;
- Compound 24: 2-(2-chloro-5-thiazolyl)-2-imidazoline;
- Compound 25: 2-(3-quinolyl)-1,4,5,6-tetrahydropyrimidine;
- Compound 26: 2-(3-furanyl)methyl-2-imidazoline;
- Compound 27: 1-(6-chloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 28: 2-(3,5-dimethyl-4-isoxazolyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 29: 2-(3,5-dimethyl-4-isoxazolyl)methyl-2-imidazoline;
- Compound 30: 2-(3-thienyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 31: 2-(3-thienyl)methyl-2-imidazoline;
- Compound 32: 2-methyl-5-(3-pyridyl)-2-imidazoline;
- Compound 33: 5-(3-pyridyl)-2-imidazoline;

- Compound 34: 1,2-bis[(6-chloro-3-pyridyl)methyl]-1,4,5,6-tetrahydropyrimidine;
- Compound 35: 1-(6-chloro-3-pyridyl)methyl-2-(3-pyridyl)-2-imidazoline;
- 5 Compound 36: 2-(5,6-dichloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 37: 2-(6-chloro-3-pyridyl)methyl-5-phenyl-1,4,5,6-tetrahydropyrimidine;
- Compound 38: 2-(4-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- 10 Compound 39: 2-(2-chloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 40: 2-(2,6-dichloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 41: 2-[2-(6-chloro-3-pyridyl)ethyl]-1,4,5,6-tetrahydropyrimidine;
- 15 Compound 42: 2-[2-(6-chloro-3-pyridyl)ethyl]-2-imidazoline;
- Compound 43: 2-(6-methyl-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 44: 1,2-bis[(6-chloro-3-pyridyl)methyl]-2-imidazoline;
- 20 Compound 45: 2-(6-methyl-3-pyridyl)methyl-2-imidazoline;
- Compound 46: 2-(6-ethoxy-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 47: 2-(6-ethoxy-3-pyridyl)methyl-2-imidazoline;
- Compound 48: 2-(6-fluoro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- 25 Compound 49: 2-(5,6-dichloro-3-pyridyl)methyl-2-imidazoline;
- Compound 50: 2-(6-chloro-3-pyridyl)methyl-5,5-dimethyl-1,4,5,6-tetrahydropyrimidine;
- Compound 51: 2-(2-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- 30 Compound 52: 1-(5,6-dichloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 53: 2-(5,6-dichloro-3-pyridyl)methyl-1-methyl-2-

imidazoline;

Compound 54: 2-(6-chloro-3-pyridyl)methyl-4-methyl-1,4,5,6-tetrahydropyrimidine;

5 Compound 55: 1-[2-(6-chloro-3-pyridyl)ethyl]-1,4,5,6-tetrahydropyrimidine;

Compound 56: 1-(3-pyridazinyl)methyl-1,4,5,6-tetrahydropyrimidine;

Compound 57: 1-(6-methyl-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;

10 Compound 58: 1-(3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;

Compound 59: 3-(6-chloro-3-pyridyl)methyl-1,4,5,6-tetrahydro-1,2,4-triazine;

Compound 60: 2-[1-(6-chloro-3-pyridyl)ethyl]-1,4,5,6-tetrahydropyrimidine;

15 Compound 61: 1-(2-chloro-5-thiazolyl)methyl-1,4,5,6-tetrahydropyrimidine;

Compound 62: 1-[2-(6-chloro-3-pyridyl)ethyl]-2-methyl-2-imidazoline;

20 Compound 63: 1-[2-(6-chloro-3-pyridyl)ethyl]-4,4-dimethyl-2-imidazoline;

Compound 64: 2-(2-chloro-5-thiazolyl)methyl-1,4,5,6-tetrahydropyrimidine;

Compound 65: 2-(2-chloro-5-thiazolyl)methyl-2-imidazoline;

Compound 66: 2-(5-pyrimidyl)methyl-1,4,5,6-tetrahydropyrimidine;

25 Compound 67: 2-(5-pyrimidyl)methyl-2-imidazoline;

Compound 68: 2-(5-methyl-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine.

30 The cyclic amidine compounds represented by the formula (I) of the present invention can be prepared in accordance with the various synthetic processes such as following Process 1 to 3.


In the following reaction schemes, the groups A<sup>1</sup>, A<sup>2</sup> and X



have the same meanings mentioned above.

**Process 1:**

In accordance with the following reaction scheme, the  
5 compound (I) of the present invention can be obtained by the  
condensation reaction of the compound of the formula (II) with  
the compound of the formula (III).



10 wherein, "Y" is  $-\text{COOQ}^1$ ,  $-\text{CONQ}^2\text{Q}^3$ ,  $-\text{C}(\text{OQ}^4)_3$ ,  $-\text{C}(\text{OQ}^5)=\text{NH}$  or  $-\text{CN}$  (in which  $\text{Q}^1$ ,  $\text{Q}^2$ ,  $\text{Q}^3$ ,  $\text{Q}^4$  and  $\text{Q}^5$  are  $\text{C}_1$ - $\text{C}_4$  lower alkyl); that is, the compound (III) represented by "A<sup>2</sup>-Y" is carboxylic acid derivative such as ester, amide, orthoester, iminoether or nitrile.

15

The compounds (II) and (III) to be used in this reaction can be commercially available or can be easily prepared from known compounds by using common methods.

The reaction of the compound (II) with the compound (III) to obtain the compound (I) can usually be carried out without solvent or in an appropriate solvent such as hydrocarbon solvent, alcohol solvent and ether solvent or the mixture thereof in the presence of acid, a reagent containing sulfur atom or an aluminum reagent if necessary, under the temperature ranging from room temperature to 300°C. The examples of acid include hydrogen chloride, p-toluenesulfonic acid and the like, and the reagent containing sulfur atom may include sulfur, hydrogen sulfide, carbon disulfide, phosphorus pentasulfide and the like.


The examples of the hydrocarbon solvent may include

aromatic hydrocarbon such as benzene, toluene and the like, or aliphatic hydrocarbon such as pentane, hexane and the like. The alcohol solvent includes methanol, ethanol, propanol, 2-propanol, 2-methyl-2-propanol ethylene glycol, diethylene glycol and the like. The examples of ether solvent may include diethyl ether, dimethoxyethane, tetrahydrofuran, 1,4-dioxane and the like.

Examples of the aluminum reagent to be used in the reaction may include trimethylaluminum, triethylaluminum, dimethylaluminum chloride, diethylaluminum chloride, ethylaluminum dichloride and the like.

#### Process 2:

The compound (I) can be obtained by the reaction of the compound (IV) with the compound (V) in accordance with the following reaction scheme.



wherein, "Z" is leaving group which accelerates the reaction with nitrogen atoms of cyclic amidine compound, such as halogen atom, p-toluenesulfonyloxy, methanesulfonyloxy, trifluoromethanesulfonyloxy, acyloxy, substituted acyloxy groups and so on.

The compounds (IV) and (V) to be used in this reaction can be commercially available or can be easily prepared from known compounds by using common methods.

The reaction of the compound (IV) with the compound (V) to obtain the compound (I) can be usually carried out in an appropriate solvent such as alcohol solvent, ketone solvent, nitrile solvent, ester solvent, amide solvent, hydrocarbon


solvent and ether solvent or the mixture thereof in the presence of an organic base or an inorganic base if necessary, under the temperature ranging from -20°C to the refluxing temperature of the solvent to be used.

5           The examples of alcohol solvent include methanol, ethanol, propanol, 2-propanol, 2-methyl-2-propanol and the like. The ketone solvent may include acetone, methyl ethyl ketone and the like. The nitrile solvent may include acetonitrile, propionitrile and so on, and the ester solvent may be ethyl  
10   acetate. The examples of amide solvent include N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidone, hexamethylphosphoramide and the like. The hydrocarbon solvent may include aromatic hydrocarbon such as benzene, toluene and the like, or aliphatic hydrocarbon such as pentane, hexane and the  
15   like. The examples of ether solvent may include diethyl ether, dimethoxyethane, tetrahydrofuran, 1,4-dioxane and the like.

Examples of the organic base to be used in the reaction may include triethylamine, collidine, lutidine, potassium tert-butoxide, sodium amide, lithium diisopropylamide, potassium  
20   bis(trimethylsilyl)amide and the like, and the inorganic base may include potassium carbonate, sodium carbonate, sodium hydrogencarbonate, sodium hydroxide, potassium hydroxide, sodium hydride, lithium hydride and the like.

25   Process 3:

The compound (I) can be obtained from the compound (VI) by the dehydrating cyclization of the compound (VI) in accordance with the following reaction scheme.



The compound (VI) to be used in this reaction can be prepared in accordance with the known method in this field.

This reaction can generally be carried out without solvent or in an appropriate solvent such as hydrocarbon solvent, 5 halogenated hydrocarbon solvent and ether solvent, or in the mixture solvent thereof, in the presence of a dehydrating reagent if necessary, at the temperature ranging from -50°C to 200°C, preferably from room temperature to 120°C.

The examples of hydrocarbon solvent may include aromatic 10 hydrocarbon such as benzene, toluene and the like, or aliphatic hydrocarbon such as pentane, hexane and the like. The examples of halogenated hydrocarbon solvent may include dichloromethane, chloroform, 1,2-dichloroethane and the like. The ether solvent may include diethyl ether, dimethoxyethane, tetrahydrofuran, 1,4- 15 dioxane and the like. The examples of the dehydrating reagent include thionyl chloride, sulfuryl chloride, phosphorus oxychloride, phosphorus trichloride, phosphorus pentachloride, p-toluenesulfonyl chloride, methanesulfonyl chloride, phosgene, diethyl azodicarboxylate, dicyclohexylcarbodiimide and the like.

20

The compound of formula (I) of the present invention thus obtained can be converted to a pharmaceutically acceptable salt with various kinds of the organic or inorganic acids mentioned above, if necessary. Furthermore, the compound (I) of the 25 present invention can also be purified by the conventional manner, such as recrystallization, column chromatography and the like.

When the compounds of the formula (I) of the present invention exist in the isomer forms, each isomer *per se* is separated from each other by the conventional manner. Therefore, 30 it is understood that each isomers *per se*, as well as the isomeric mixture, shall be included in the compounds of the present invention.

The compounds of formula (I) of the present invention bind selectively to nicotinic acetylcholine receptors in central nervous systems, and activate said receptors as agonists or modulators. Therefore, these compounds are useful as medicaments  
5 for preventing or treating various diseases, such as dementia, senile dementia, presenile dementia, Alzheimer's disease, Parkinson's disease, cerebrovascular dementia, AIDS-related dementia, dementia in Down's syndrome, Tourette's syndrome, neurosis during chronic cerebral infarction stage, cerebral  
10 dysfunction caused by cerebral injury, anxiety, schizophrenia, depression, Huntington's disease, pain and so on.

The compounds of formula (I) or a pharmaceutically acceptable salt thereof according to the present invention may be administered in the form of oral or parenteral formulations. The  
15 formulations for oral administration may include for example, tablets, capsules, granules, fine powders, syrups or the like; the formulations for parenteral administration may include, for example, injectable solutions or suspensions with distilled water for injection or other pharmaceutically acceptable solution,  
20 patches for transdermal application, sprays for nasally administration, depositories or the like.

These formulations may be formed by mixing with pharmaceutically acceptable carrier, excipient, sweetener, stabilizer and so on by the conventional procedures known *per se*  
25 to those skilled in the art in the field of pharmaceutical formulations.

Examples of pharmaceutically acceptable carrier or excipient include polyvinyl pyrrolidone, gum arabic, gelatin, sorbit, cyclodextrin, magnesium stearate, talc, polyethylene  
30 glycol, polyvinyl alcohol, silica, lactose, crystalline cellulose, sugar, starch, calcium phosphate, vegetable oil, carboxymethyl cellulose, hydroxypropyl cellulose, sodium lauryl sulfate, water,

ethanol, glycerol, mannitol, syrup and the like.

The solutions for injection may be isotonic solution containing glucose and the like, and these solutions can be further contain an appropriate solubilizer such as polyethylene glycol or the like, buffer, stabilizer, preservative, antioxidant and so on.

These formulations can be administered to the human being and other mammalian animals, and the preferable administration route may include oral route, transdermic route, nasal route, rectal route, topical route or the like.

The administration dose may vary in a wide range with ages, weights, condition of patients, routes of administration or the like, and a usual recommended daily dose to adult patients for oral administration is within the range of approximately 0.001-1,000 mg/kg per body weight, preferably 0.01-100 mg/kg per body weight, and more preferably 0.1-10 mg/kg per body weight.

In the case of parenteral administration such as intravenous injections, a usual recommended daily dose is within the range of approximately 0.00001-10 mg/kg per body weight, preferably 0.0001-1 mg/kg per body weight, and more preferably 0.001-0.1 mg/kg per body weight, once or in three times per day.

The methods for evaluating the binding capabilities of the compounds at nicotinic acetylcholine receptors are different by subtypes of receptors. The binding capabilities of the compounds at  $\alpha 4 \beta 2$  nicotinic acetylcholine receptors are examined using rat brain membrane obtained from whole homogenized brain, and determining the inhibiting rate of the compounds against [ $^3$ H]-cytisine binding to said brain membrane. Furthermore, the binding capabilities of the compounds at  $\alpha 1 \beta 1 \gamma \delta$  nicotinic acetylcholine receptors are examined using homogenized rat muscle, and determining the inhibiting rate of the compounds against

[<sup>3</sup>H]- $\alpha$ -bungarotoxin binding to said muscle homogenate.

The agonist effect in human  $\alpha 4\beta 2$  subtype of nicotinic acetylcholine receptors are examined by using human nicotinic acetylcholine receptors prepared in oocytes of *Xenopus laevis*,  
5 which is injected with cRNA from the corresponding cloning cDNA of human  $\alpha 4$  and  $\beta 2$  subunits of nicotinic acetylcholine receptors, and to measure the expression of the electric response by adding the test compounds to perfusion solution by means of membrane potential holding method.

10

Examples:

The present invention is illustrated in more detail by way of the following examples.

15 Example 1: Synthesis by the Process 1

2-(6-chloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine

[Compound 8]

To a stirred solution of 20 ml of toluene were added 3.75 ml of 1M trimethylaluminum/hexane solution and 315  $\mu$ l (3.77 mmol)  
20 of trimethylenediamine under argon atmosphere at room temperature, and to this mixture was further added 500 mg (2.5 mmol) of ethyl (6-chloro-3-pyridyl)acetate in toluene solution. The mixture was stirred for 22 hours at 100°C under refluxing. After cooling the reaction mixture to room temperature, 5 ml of chloroform, 5 ml of  
25 methanol and 1 ml of water were added. Then precipitated gel was removed off by filtration and washed with a mixture of chloroform and methanol (9:1), and the filtrate was concentrated under reduced pressure. The resulting residue was purified by aminopropyl-coated silica gel (Chromatorex NH-type; Fuji Silysia  
30 Chemical Ltd.) column chromatography (eluent; dichloromethane : ethyl acetate = 30:1, then dichloromethane : methanol = 50:1) to give 442 mg (yield; 84.4%) of 2-(6-chloro-3-pyridyl)methyl-

1,4,5,6-tetrahydropyrimidine as crystalline. This product was dissolved in methanol and to this solution was added 245 mg (2.11 mmol) of fumaric acid, and the mixture was concentrated under reduced pressure. The resulting oily residue was treated with acetonitrile to give crystalline. The crystalline was collected by filtration and dried in vacuum to give 643 mg of fumarate of the title Compound 8.

The following compounds were synthesized in accordance with the same procedures as described in Example 1.

Compound 1: 2-(6-chloro-3-pyridyl)-2-imidazoline;

Compound 2: 2-(6-chloro-3-pyridyl)-1,4,5,6-tetrahydropyrimidine;

Compound 3: 2-(6-chloro-3-pyridyl)-1-methyl-2-imidazoline;

Compound 4: 2-(6-chloro-3-pyridyl)-1-methyl-1,4,5,6-tetrahydro-pyrimidine;

Compound 6: 2-(6-chloro-3-pyridyl)imidazole;

Compound 7: 2-(6-chloro-3-pyridyl)methyl-2-imidazoline;

Compound 9: 2-(6-chloro-3-pyridyl)methyl-1-methyl-2-imidazoline;

Compound 10: 2-(6-chloro-3-pyridyl)methyl-1-methyl-1,4,5,6-tetrahydro-pyrimidine;

Compound 13: 2-(tetrahydrofuran-3-yl)-1,4,5,6-tetrahydro-pyrimidine;

Compound 14: 2-(tetrahydrofuran-3-yl)-2-imidazoline;

Compound 15: 2-(tetrahydrofuran-3-yl)methyl-1,4,5,6-tetrahydro-pyrimidine;

Compound 16: 2-(5-bromo-3-pyridyl)methyl-1,4,5,6-tetrahydro-pyrimidine;

Compound 17: 2-(5-bromo-3-pyridyl)methyl-2-imidazoline;

Compound 18: 2-(3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;

Compound 19: 2-(3-pyridyl)methyl-2-imidazoline;

Compound 20: 2-(3-aminophenyl)-1,4,5,6-tetrahydropyrimidine;

Compound 21: 2-(3-quinolyl)methyl-1,4,5,6-tetrahydropyrimidine;



- Compound 22: 2-(2-chloro-5-thiazolyl)-1,4,5,6-tetrahydro-pyrimidine;
- Compound 23: 2-(3-quinolyl)methyl-2-imidazoline;
- Compound 24: 2-(2-chloro-5-thiazolyl)-2-imidazoline;
- 5 Compound 25: 2-(3-quinolyl)-1,4,5,6-tetrahydropyrimidine;
- Compound 26: 2-(3-furanyl)methyl-2-imidazoline;
- Compound 28: 2-(3,5-dimethyl-4-isoxazolyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 29: 2-(3,5-dimethyl-4-isoxazolyl)methyl-2-imidazoline;
- 10 Compound 30: 2-(3-thienyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 31: 2-(3-thienyl)methyl-2-imidazoline;
- Compound 33: 5-(3-pyridyl)-2-imidazoline;
- Compound 36: 2-(5,6-dichloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- 15 Compound 37: 2-(6-chloro-3-pyridyl)methyl-5-phenyl-1,4,5,6-tetrahydropyrimidine;
- Compound 38: 2-(4-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 39: 2-(2-chloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- 20 Compound 40: 2-(2,6-dichloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 41: 2-[2-(6-chloro-3-pyridyl)ethyl]-1,4,5,6-tetrahydropyrimidine;
- Compound 42: 2-[2-(6-chloro-3-pyridyl)ethyl]-2-imidazoline;
- 25 Compound 43: 2-(6-methyl-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- Compound 45: 2-(6-methyl-3-pyridyl)methyl-2-imidazoline;
- Compound 46: 2-(6-ethoxy-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- 30 Compound 47: 2-(6-ethoxy-3-pyridyl)methyl-2-imidazoline;
- Compound 48: 2-(6-fluoro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;

Compound 49: 2-(5,6-dichloro-3-pyridyl)methyl-2-imidazoline;

Compound 50: 2-(6-chloro-3-pyridyl)methyl-5,5-dimethyl-1,4,5,6-tetrahydropyrimidine;

Compound 51: 2-(2-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;

5 Compound 53: 2-(5,6-dichloro-3-pyridyl)methyl-1-methyl-2-imidazoline;

Compound 54: 2-(6-chloro-3-pyridyl)methyl-4-methyl-1,4,5,6-tetrahydropyrimidine;

10 Compound 59: 3-(6-chloro-3-pyridyl)methyl-1,4,5,6-tetrahydro-1,2,4-triazine;

Compound 60: 2-[1-(6-chloro-3-pyridyl)ethyl]-1,4,5,6-tetrahydropyrimidine;

Compound 61: 1-(2-chloro-5-thiazolyl)methyl-1,4,5,6-tetrahydropyrimidine;

15 Compound 62: 1-[2-(6-chloro-3-pyridyl)ethyl]-2-methyl-2-imidazoline;

Compound 63: 1-[2-(6-chloro-3-pyridyl)ethyl]-4,4-dimethyl-2-imidazoline;

20 Compound 64: 2-(2-chloro-5-thiazolyl)methyl-1,4,5,6-tetrahydropyrimidine;

Compound 65: 2-(2-chloro-5-thiazolyl)methyl-2-imidazoline;

Compound 66: 2-(5-pyrimidyl)methyl-1,4,5,6-tetrahydropyrimidine;

Compound 67: 2-(5-pyrimidyl)methyl-2-imidazoline;

25 Compound 68: 2-(5-methyl-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine.

#### Example 2: Synthesis by the Process 2

##### 1-(6-chloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine

##### [Compound 27]

30 To an ice-cooled solution of 384 mg (4.6 mmol) of 1,4,5,6-tetrahydropyrimidine in 5 ml of acetonitrile was added 619 mg (3 mmol) of 5-bromomethyl-2-chloropyridine, and the mixture was

stirred for 15 minutes. After removal of solvent under reduced pressure, 6 ml of the solution of 0.5N potassium hydroxide in ethanol was added to the residue. The insoluble matter was removed off by filtration, and the filtrate was concentrated under reduced pressure. The resulting residue was dissolved in toluene, and the solvent was removed again under reduced pressure. The resulting residue was purified by aminopropyl-coated silica gel (Chromatorex NH-type; Fuji Silysia Chemical Ltd.) column chromatography (eluent; dichloromethane : methanol = 40:1) to give 221 mg (yield; 35.2%) of 1-(6-chloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine as colorless oil. This product was dissolved in methanol and to this solution was added 122 mg (1.05 mmol) of fumaric acid, and the mixture was concentrated under reduced pressure. The resulting residue was treated with acetonitrile to give crystalline. The crystalline was collected by filtration and dried in vacuum to give 308 mg of fumarate of the title Compound 27.

The following compounds were synthesized in accordance with the same procedures as described in Example 2.

Compound 5: 1-(6-chloro-3-pyridyl)methylimidazole;

Compound 10: 2-(6-chloro-3-pyridyl)methyl-1-methyl-1,4,5,6-tetrahydropyrimidine;

Compound 11: 1-(6-chloro-3-pyridyl)methyl-2-methyl-2-imidazoline;

Compound 34: 1,2-bis[(6-chloro-3-pyridyl)methyl]-1,4,5,6-tetrahydropyrimidine;

Compound 35: 1-(6-chloro-3-pyridyl)methyl-2-(3-pyridyl)-2-imidazoline;

Compound 44: 1,2-bis[(6-chloro-3-pyridyl)methyl]-2-imidazoline;

Compound 52: 1-(5,6-dichloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;

Compound 55: 1-[2-(6-chloro-3-pyridyl)ethyl]-1,4,5,6-tetrahydro-

pyrimidine;

Compound 56: 1-(3-pyridazinyl)methyl-1,4,5,6-tetrahydro-pyrimidine;

5                      Compound 57: 1-(6-methyl-3-pyridyl)methyl-1,4,5,6-tetrahydro-pyrimidine;

Compound 58: 1-(3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;

Compound 61: 1-(2-chloro-5-thiazolyl)methyl-1,4,5,6-tetrahydro-pyrimidine;

10                    Compound 62: 1-[2-(6-chloro-3-pyridyl)ethyl]-2-methyl-2-imidazoline;

Compound 63: 1-[2-(6-chloro-3-pyridyl)ethyl]-4,4-dimethyl-2-imidazoline.

Example 3: Synthesis by the Process 3

15    2-Methyl-5-(3-pyridyl)-2-imidazoline [Compound 32]

269 mg (1 mmol) of oxalate of N-[2-amino-1-(3-pyridyl)ethyl]acetamide was dissolved in 5 ml of phosphorus oxychloride, and this mixture was heated for 1.5 hours at 100°C under stirring. After cooling the reaction mixture to room  
20 temperature, phosphorus oxychloride was removed off under reduced pressure. The resulting residue was treated with ice, and 1N sodium hydroxide aqueous solution was added to adjust the pH of the solution to 7, then, the mixture was concentrated under reduced pressure. The resulting residue was treated with ethanol  
25 and the insoluble matter was removed off by filtration, and the filtrate was concentrated under reduced pressure. The resulting residue was purified by aminopropyl-coated silica gel (Chromatorex NH-type; Fuji Silysia Chemical Ltd.) column chromatography (eluent; chloroform) to give 22 mg (yield; 13.6%)  
30 of 2-methyl-5-(3-pyridyl)-2-imidazoline as brownish oil. This product was dissolved in methanol and to this solution was added 15 mg (0.13 mmol) of fumaric acid, and the mixture was

concentrated under reduced pressure. The resulting oily residue was treated with a mixture of t-butanol and acetone to give crystalline. The crystalline was collected by filtration and dried in vacuum to give 17 mg of fumarate of the title Compound

5 32.

The physicochemical data of the Compounds 1 to 68 obtained by the above-mentioned examples are summarized in the following Table 1 to Table 14.

TABLE 1:






No.	Chemical Structure	Salt	Properties m.p. (°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR (DMSO-d <sub>6</sub> )
1		fumarate	colorless cryst. 170-175°C acetonitrile	m/z 182 = (M+H) <sup>+</sup> C <sub>8</sub> H <sub>8</sub> ClN <sub>3</sub>	8.87 (d, J=2.4Hz, 1H), 8.29 (dd, J=2.4, 8.4Hz, 1H), 7.70 (d, J=8.4Hz, 1H), 6.56 (s, 2H), 3.78 (s, 4H)
2		fumarate	colorless cryst. 162-168°C methanol /acetonitrile	m/z 196 = (M+H) <sup>+</sup> C <sub>9</sub> H <sub>10</sub> ClN <sub>3</sub>	8.79 (d, J=2.5Hz, 1H), 8.24 (dd, J=2.5, 8.3Hz, 1H), 7.74 (d, J=8.3Hz, 1H), 6.40 (s, 2H), 3.49 (t, J=5.7Hz, 4H), 1.94 (m, 2H)
3		fumarate	milky white cryst. 117-120°C ether	m/z 196 = (M+H) <sup>+</sup> C <sub>9</sub> H <sub>10</sub> ClN <sub>3</sub>	8.65 (d, J=2.4Hz, 1H), 8.09 (dd, J=2.4, 8.2Hz, 1H), 7.71 (d, J=8.2Hz, 1H), 6.53 (s, 2H), 3.84 (m, 2H), 3.70 (m, 2H), 2.89 (s, 3H)
4		oxalate	colorless oil	m/z 210 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>12</sub> ClN <sub>3</sub>	10.26 (br, 1H) 8.66 (d, J=1.8Hz, 1H), 8.13 (dd, J=1.8, 8.3Hz, 1H), 7.80 (d, J=8.3Hz, 1H), 3.57 (t, J=5.6Hz, 2H), 3.43 (t, J=5.3Hz, 2H), 2.98 (s, 3H), 2.08 (m, 2H)
5		fumarate	colorless cryst. 123-124°C acetonitrile	m/z 194 = (M+H) <sup>+</sup> C <sub>9</sub> H <sub>8</sub> ClN <sub>3</sub>	8.39 (d, J=2.4Hz, 1H), 7.81 (d, J=4.6Hz, 1H), 7.73 (dd, J=2.4, 8.2Hz, 1H), 7.52 (d, J=8.2Hz, 1H), 7.24 (s, 1H), 6.94 (br, 1H), 6.63 (s, 2H), 5.26 (s, 2H)

TABLE 2:






No.	Chemical Structure	Salt	Properties m.p. (°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR(DMSO-d <sub>6</sub> )
6		fumarate	colorless cryst. 173-186°C acetonitrile	m/z 180 = (M+H) <sup>+</sup> C <sub>8</sub> H <sub>8</sub> ClN <sub>3</sub>	13.0 (br, 3H), 8.94 (d, J=2.5Hz, 1H), 8.30 (dd, J=2.5, 8.3Hz, 1H), 7.60 (d, J=8.3Hz, 1H), 7.23 (s, 2H), 6.63 (s, 2H)
7		fumarate	colorless cryst. 139-142°C acetonitrile	m/z 196 = (M+H) <sup>+</sup> C <sub>8</sub> H <sub>10</sub> ClN <sub>3</sub>	8.42 (d, J=2.5Hz, 1H), 7.87 (dd, J=2.5, 8.2Hz, 1H), 7.52 (d, J=8.2Hz, 1H), 6.47 (s, 2H), 3.93 (s, 2H), 3.73 (s, 4H)
8		fumarate	colorless cryst. 167-172°C acetonitrile	m/z 210 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>12</sub> ClN <sub>3</sub>	8.46 (d, J=2.5Hz, 1H), 7.92 (dd, J=2.5, 8.3Hz, 1H), 7.52 (d, J=8.3Hz, 1H), 6.45 (s, 2H), 3.87 (s, 2H), 3.32 (t, J=5.7Hz, 4H), 1.81 (m, 2H)
9		fumarate	colorless cryst. 123-126°C acetonitrile	m/z 210 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>12</sub> ClN <sub>3</sub>	8.43 (br, 1H), 7.86 (dd, J=2.3, 8.2Hz, 1H), 7.54 (d, J=8.2Hz, 1H), 6.48 (s, 2H), 4.06 (s, 2H), 3.76 (m, 4H), 3.00 (s, 3H)
10		oxalate	colorless cryst. 85-89°C acetone	m/z 224 = (M+H) <sup>+</sup> C <sub>11</sub> H <sub>14</sub> ClN <sub>3</sub>	8.42 (d, J=2.4Hz, 1H), 7.84 (dd, J=2.4, 8.2Hz, 1H), 7.55 (d, J=8.2Hz, 1H), 4.07 (s, 2H), 3.44 (t, J=5.7Hz, 2H), 3.35 (t, J=5.7Hz, 2H), 3.06 (s, 3H), 1.95 (m, 2H)

TABLE 3:






No.	Chemical Structure	Salt	Properties m.p. (°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR(DMSO-d <sub>6</sub> )
11		fumarate	colorless cryst. 165-171°C acetonitrile	m/z 210 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>12</sub> ClN <sub>3</sub>	8.45 (d, J=2.5Hz, 1H), 7.89 (dd, J=2.5, 8.2Hz, 1H), 7.57 (d, J=8.2Hz, 1H), 6.46 (s, 2H), 4.63 (s, 2H), 3.73 (m, 2H), 3.63 (m, 2H), 2.32 (s, 3H)
12		fumarate	colorless cryst. 166-168°C acetonitrile	m/z 224 = (M+H) <sup>+</sup> C <sub>11</sub> H <sub>14</sub> ClN <sub>3</sub>	8.41 (d, J=2.5Hz, 1H), 7.95 (s, 1H), 7.86 (dd, J=2.5, 8.2Hz, 1H), 7.56 (d, J=8.2Hz, 1H), 6.49 (s, 2H), 4.57 (s, 2H), 3.17 (s, 2H), 1.24 (s, 6H)
13		fumarate	pale yellow cryst. 54-57°C acetone	m/z 155 = (M+H) <sup>+</sup> C <sub>8</sub> H <sub>14</sub> N <sub>2</sub> O	9.9 (br, 1H), 6.43 (s, 2H), 3.88 (m, 2H), 3.72 (m, 2H), 3.31 (t, J=5.7Hz, 4H), 3.29 (m, 1H), 2.21 (m, 1H), 2.04 (m, 1H), 1.84 (quintet, J=5.7Hz, 2H)
14		fumarate	colorless cryst. 103-105°C acetone	m/z 141 = (M+H) <sup>+</sup> C <sub>7</sub> H <sub>12</sub> N <sub>2</sub> O	6.43 (s, 2H), 3.86 (m, 2H), 3.73 (s, 4H), 3.72 (m, 2H), 3.35 (m, 1H), 2.19 (m, 1H), 2.06 (m, 1H)
15		oxalate	colorless cryst. 187-190°C acetone	m/z 169 = (M+H) <sup>+</sup> C <sub>9</sub> H <sub>16</sub> N <sub>2</sub> O	9.71 (br, 2H), 3.74 (m, 2H), 3.64 (m, 1H), 3.32 (m, 4H), 2.44 (m, 4H), 1.99 (m, 1H), 1.84 (m, 2H), 1.54 (m, 1H)



TABLE 4:






No.	Chemical Structure	Salt	Properties m.p. (°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR (DMSO-d <sub>6</sub> )
16		fumarate	colorless cryst. 155-159°C acetone	m/z 254 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>12</sub> BrN <sub>3</sub>	8.66 (d, J=1.6Hz, 1H), 8.62 (d, J=1.6Hz, 1H), 8.16 (s, 1H), 6.39 (s, 2H), 3.87 (s, 2H), 3.33 (m, 4H), 1.81 (m, 2H)
17		fumarate	colorless cryst. 150-154°C acetone	m/z 242 = (M+H) <sup>+</sup> C <sub>9</sub> H <sub>10</sub> BrN <sub>3</sub>	8.63 (s, 1H), 8.53 (s, 1H), 8.05 (s, 1H), 6.44 (s, 2H), 3.78 (s, 2H), 3.65 (s, 4H)
18		fumarate	colorless cryst. 120-124°C ethanol /acetone	m/z 176 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>13</sub> N <sub>3</sub>	10.77 (2H, br), 8.62 (1H, s), 8.51 (d, J=4.8Hz, 1H), 7.85 (d, J=7.6Hz, 1H), 7.39 (dd, J=4.8, 7.6Hz, 1H), 6.42 (s, 2H), 3.86 (s, 2H), 3.33 (m, 4H), 1.81 (m, 2H)
19		fumarate	colorless cryst. 134-135°C acetone	m/z 162 = (M+H) <sup>+</sup> C <sub>8</sub> H <sub>11</sub> N <sub>3</sub>	8.57 (d, J=2.0Hz, 1H), 8.51 (dd, J=2.0, 4.7Hz, 1H), 7.78 (d, J=7.8Hz, 1H), 7.39 (dd, J=4.7, 7.8Hz, 1H), 6.46 (s, 2H), 3.85 (s, 2H), 3.72 (s, 4H)
20		fumarate	colorless cryst. 192-195°C acetone	m/z 176 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>13</sub> N <sub>3</sub>	7.21 (m, 1H), 6.85 (s, 1H), 6.81 (m, 2H), 6.37 (s, 2H), 5.54 (br, 2H), 3.45 (m, 4H), 1.95 (m, 2H)

TABLE 5:











No.	Chemical Structure	Salt	Properties m.p.(°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR(DMSO-d <sub>6</sub> )
21		fumarate	colorless cryst. 168-171°C acetone	m/z 226 = (M+H) <sup>+</sup> C <sub>14</sub> H <sub>15</sub> N <sub>3</sub>	8.94 (s, 1H), 8.38 (s, 1H), 8.03 (d, J=8.4Hz, 1H), 7.94 (d, J=8.1Hz, 1H), 7.77 (m, 1H), 7.64 (m, 1H), 6.42 (s, 2H), 4.05 (s, 2H), 3.34 (m, 4H), 1.83 (m, 2H)
22		fumarate	colorless cryst. 159-160°C acetone	m/z 202 = (M+H) <sup>+</sup> C <sub>7</sub> H <sub>8</sub> ClN <sub>3</sub> S	8.03 (s, 1H), 6.56 (s, 2H), 3.34(m, 4H), 1.76 (m, 2H)
23		fumarate	colorless cryst. 175-177°C acetone	m/z 212 = (M+H) <sup>+</sup> C <sub>13</sub> H <sub>13</sub> N <sub>3</sub>	8.88 (s, 1H), 8.31 (s, 1H), 8.03 (d, J=8.4Hz, 1H), 7.96 (d, J=8.1Hz, 1H), 7.78 (m, 1H), 7.64 (m, 1H), 6.47 (s, 2H), 4.06 (s, 2H), 3.75 (s, 4H)
24		fumarate	pale yellow cryst. 157-158°C acetone	m/z 188 = (M+H) <sup>+</sup> C <sub>6</sub> H <sub>6</sub> ClN <sub>3</sub> S	8.02 (s, 1H), 6.62 (s, 2H), 3.62(s, 4H)
25		fumarate	colorless cryst. 188-193°C acetone	m/z 212 = (M+H) <sup>+</sup> C <sub>13</sub> H <sub>13</sub> N <sub>3</sub>	9.16 (d, J=2.2Hz, 1H), 8.82 (d, J=2.2Hz, 1H), 8.13 (m, 2H), 7.95 (m, 1H), 7.76 (m, 1H), 6.38 (s, 2H), 3.55 (m, 4H), 2.00 (m, 2H)

TABLE 6:

No.	Chemical Structure	Salt	Properties m.p. (°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR (DMSO-d <sub>6</sub> )
26		fumarate	colorless cryst. 200-205°C acetone	m/z 151 = (M+H) <sup>+</sup> C <sub>8</sub> H <sub>10</sub> N <sub>2</sub> O	7.66 (s, 1H), 7.64 (s, 1H), 6.50 (s, 1H), 6.41 (s, 2H), 3.74 (s, 4H), 3.69 (s, 2H)
27		fumarate	colorless cryst. 126-129°C acetonitrile	m/z 210 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>12</sub> ClN <sub>3</sub>	8.47 (m, 2H), 7.92 (dd, J=2.5, 8.2Hz, 1H), 7.59 (d, J=8.2Hz, 1H), 6.44 (s, 2H), 4.69 (s, 2H), 3.25 (m, 4H), 1.88 (m, 2H)
28		fumarate	colorless cryst. 188-190°C acetone	m/z 194 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>15</sub> N <sub>3</sub> O	10.37 (br, 2H), 6.39 (s, 2H), 3.68 (s, 2H), 3.32 (m, 4H), 2.34 (s, 3H), 2.14 (s, 3H), 1.83 (m, 2H)
29		fumarate	colorless cryst. 208-215°C ethanol	m/z 180 = (M+H) <sup>+</sup> C <sub>9</sub> H <sub>13</sub> N <sub>3</sub> O	6.43 (s, 2H), 3.72 (s, 4H), 3.64 (s, 2H), 2.34 (s, 3H), 2.14 (s, 3H)
30		fumarate	colorless cryst. 85-90°C acetone	m/z 181 = (M+H) <sup>+</sup> C <sub>9</sub> H <sub>12</sub> N <sub>2</sub> S	7.55 (d, J=4.8Hz, 1H), 7.46 (s, 1H), 7.13 (d, J=4.8Hz, 1H), 6.40 (s, 2H), 3.78 (s, 2H), 3.33 (m, 4H), 1.83 (m, 2H)

**TABLE 7:**

No.	Chemical Structure	Salt	Properties m.p. (°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR(DMSO-d <sub>6</sub> )
31		fumarate	colorless cryst. 150–153°C	m/z 167 = (M+H) <sup>+</sup> C <sub>8</sub> H <sub>10</sub> N <sub>2</sub> S	7.55 (d, J=4.8Hz, 1H), 7.43 (s, 1H), 7.11 (d, J=4.8Hz, 1H), 6.43 (s, 2H), 3.83 (s, 2H), 3.75 (s, 4H)
32		fumarate	pale brown cryst. 130–132°C t-butanol /acetone	m/z 162 = (M+H) <sup>+</sup> C <sub>9</sub> H <sub>11</sub> N <sub>3</sub>	8.60 (s, 1H), 8.57 (m, 1H), 7.81 (d, J=6.8Hz, 1H), 7.45 (m, 1H), 6.48 (s, 2H), 5.33 (m, 1H), 4.23 (m, 1H), 3.64 (m, 1H), 2.24 (s, 3H)
33		fumarate	colorless cryst. 148–149°C ethanol /acetone	m/z 148 = (M+H) <sup>+</sup> C <sub>8</sub> H <sub>9</sub> N <sub>3</sub>	8.56 (m, 2H), 8.14 (s, 1H), 7.75 (d, J=7.0Hz, 1H), 7.43 (m, 1H), 6.54 (s, 2H), 5.24 (m, 1H), 4.15 (m, 1H), 3.55 (m, 1H)
34		fumarate	pale brown cryst. 135–139°C	m/z 335 = (M+H) <sup>+</sup> C <sub>16</sub> H <sub>16</sub> Cl <sub>2</sub> N <sub>4</sub>	8.40 (d, J=2.3Hz, 1H), 8.20 (s, 1H), 7.84 (dd, J=2.3, 8.3Hz, 1H), 7.64 (d, J=8.2Hz, 1H), 7.47 (m, 2H), 6.47 (s, 2H), 4.74 (s, 2H), 4.23 (s, 2H), 3.42 (t, J=5.4Hz, 2H), 3.34 (t, J=5.3Hz, 2H), 1.96 (m, 2H)
35		fumarate	pale brown cryst. 164–166°C	m/z 273 = (M+H) <sup>+</sup> C <sub>14</sub> H <sub>13</sub> ClN <sub>4</sub>	8.76 (d, J=1.8Hz, 1H), 8.71 (dd, J=1.5, 4.8Hz, 1H), 8.34 (d, J=2.4Hz, 1H), 7.97 (ddd, J=1.5, 1.8, 7.8Hz, 1H), 7.81 (dd, J=2.4, 8.2Hz, 1H), 7.53 (dd, J=4.8, 7.8Hz, 1H), 7.52 (d, J=8.2Hz, 1H), 6.58 (s, 2H), 4.32 (s, 2H), 3.83 (t, J=10.0Hz, 2H), 3.45 (t, J=10.0Hz, 2H)

TABLE 8:






No.	Chemical Structure	Salt	Properties m.p. (°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR (DMSO-d <sub>6</sub> )
36		fumarate	colorless cryst. 198-200°C acetone	m/z 244 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>11</sub> Cl <sub>2</sub> N <sub>3</sub>	8.31 (d, J=2.1Hz, 1H), 8.01 (d, J=2.1Hz, 1H), 6.68 (s, 2H), 3.85 (s, 2H), 3.43 (m, 4H), 1.99 (m, 2H) in CD <sub>3</sub> OD
37		fumarate	colorless cryst. 163-168°C acetone	m/z 286 = (M+H) <sup>+</sup> C <sub>16</sub> H <sub>18</sub> ClN <sub>3</sub>	8.49 (d, J=2.4Hz, 1H), 7.94 (dd, J=2.4, 8.2Hz, 1H), 7.55 (d, J=8.2Hz, 1H), 7.30 (m, 5H), 6.44 (s, 2H), 3.94 (s, 2H), 3.57 (m, 2H), 3.45 (m, 2H), 3.08 (m, 1H)
38		fumarate	colorless cryst. 141-143°C acetone	m/z 176 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>13</sub> N <sub>3</sub>	8.55 (d, J=5.8Hz, 2H), 7.40 (d, J=5.8Hz, 2H), 6.48 (s, 2H), 3.84 (s, 2H), 3.34 (t, J=5.7Hz, 4H), 1.83 (m, 2H)
39		fumarate	colorless cryst. 160-161°C acetone	m/z 210 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>12</sub> ClN <sub>3</sub>	8.38 (dd, J=1.7, 4.8Hz, 1H), 7.89 (dd, J=1.7, 7.6Hz, 1H), 7.46 (dd, J=4.8, 7.6Hz, 1H), 6.35 (s, 2H), 3.97 (s, 2H), 3.35 (t, J=5.7Hz, 4H), 1.87 (m, 2H)
40		fumarate	colorless cryst. 175-177°C acetone	m/z 244 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>11</sub> Cl <sub>2</sub> N <sub>3</sub>	7.86 (d, J=8.0Hz, 1H), 7.50 (d, J=8.0Hz, 1H), 6.68 (s, 2H), 3.97 (s, 2H), 3.45 (t, J=5.7Hz, 4H), 2.02 (m, 2H) in CD <sub>3</sub> OD

TABLE 9:






No.	Chemical Structure	Salt	Properties m.p. (°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR (DMSO-d <sub>6</sub> )
41		fumarate	colorless cryst. 156–157°C acetone	m/z 224 = (M+H) <sup>+</sup> C <sub>11</sub> H <sub>14</sub> ClN <sub>3</sub>	8.28 (s, 1H), 7.74 (d, J=8.2Hz, 1H), 7.46 (d, J=8.2Hz, 1H), 6.70 (s, 2H), 3.41 (t, J=5.5Hz, 4H), 3.02 (t, J=7.6Hz, 2H), 2.73 (t, J=7.6Hz, 2H), 1.95 (m, 2H) in CD <sub>3</sub> OD
42		fumarate	colorless cryst. 148–149°C acetone	m/z 210 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>12</sub> ClN <sub>3</sub>	8.27 (s, 1H), 7.73 (d, J=8.0Hz, 1H), 7.43 (d, J=8.0Hz, 1H), 6.68 (s, 2H), 3.90 (s, 4H), 3.02 (br, 2H), 2.86 (br, 2H) in CD <sub>3</sub> OD
43		fumarate	colorless cryst. 156–158°C 2-propanol /acetone	m/z 190 = (M+H) <sup>+</sup> C <sub>11</sub> H <sub>16</sub> N <sub>3</sub>	8.46 (s, 1H), 7.71 (d, J=7.9Hz, 1H), 7.23 (d, J=7.9Hz, 1H), 6.40 (s, 2H), 3.77 (s, 2H), 3.31 (m, 4H), 2.44 (s, 3H), 1.80 (m, 2H)
44		fumarate	milky white cryst. 162–164°C 2-propanol	m/z 321 = (M+H) <sup>+</sup> C <sub>19</sub> H <sub>14</sub> Cl <sub>2</sub> N <sub>4</sub>	8.38 (d, J=2.0Hz, 1H), 8.31 (d, J=2.4Hz, 1H), 7.82 (dd, J=2.0, 8.2Hz, 1H), 7.75 (dd, J=2.4, 8.2Hz, 1H), 7.51 (d, J=8.2Hz, 1H), 7.49 (d, J=8.2Hz, 1H), 6.52 (s, 2H), 4.57 (s, 2H), 4.00 (s, 2H), 3.68 (m, 2H), 3.47 (m, 2H)
45		fumarate	colorless cryst. 165–166°C acetone	m/z 176 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>13</sub> N <sub>3</sub>	8.42 (d, J=2.2Hz, 1H), 7.66 (dd, J=2.2, 8.0Hz, 1H), 7.23 (d, J=8.0Hz, 1H), 6.44 (s, 2H), 3.82 (s, 2H), 3.72 (s, 4H), 2.44 (s, 3H)

TABLE 10:






No.	Chemical Structure	Salt	Properties m.p. (°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR (DMSO-d <sub>6</sub> )
46		fumarate	colorless cryst. 110-112°C acetone	m/z 220 = (M+H) <sup>+</sup> C <sub>12</sub> H <sub>17</sub> N <sub>3</sub> O	8.16 (d, J=2.3Hz, 1H), 7.72 (dd, J=2.3, 8.5Hz, 1H), 6.78 (d, J=8.5Hz, 1H), 6.39 (s, 2H), 4.28 (q, J=7.0Hz, 2H), 3.72 (s, 2H), 3.31 (t, J=5.7Hz, 4H), 1.80 (m, 2H), 1.30 (t, J=7.0Hz, 3H)
47		fumarate	colorless cryst. 170-171°C acetone	m/z 206 = (M+H) <sup>+</sup> C <sub>11</sub> H <sub>15</sub> N <sub>3</sub> O	8.12 (d, J=2.2Hz, 1H), 7.68 (dd, J=2.2, 8.5Hz, 1H), 6.78 (d, J=8.5Hz, 1H), 6.42 (s, 2H), 4.27 (q, J=7.0Hz, 2H), 3.76 (s, 2H), 3.72 (s, 4H), 1.30 (t, J=7.0Hz, 3H)
48		fumarate	pale yellow cryst. 136-139°C acetone	m/z 194 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>12</sub> FN <sub>3</sub>	8.27 (s, 1H), 8.03 (ddd, J=2.3, 8.2, 8.4Hz, 1H), 7.21 (dd, J=8.4, 2.7Hz, 1H), 6.39 (s, 2H), 3.84 (s, 2H), 3.32 (t, J=5.7, 4H), 1.81 (m, 2H)
49		fumarate	colorless cryst. 176-178°C acetone	m/z 230 = (M+H) <sup>+</sup> C <sub>9</sub> H <sub>9</sub> Cl <sub>2</sub> N <sub>3</sub>	8.37 (s, 1H), 8.15 (s, 1H), 6.46 (s, 2H), 3.85 (s, 2H), 3.66 (s, 4H)
50		fumarate	pale yellow cryst. 143-145°C acetone	m/z 238 = (M+H) <sup>+</sup> C <sub>12</sub> H <sub>18</sub> ClN <sub>3</sub>	8.37 (s, 1H), 7.82 (dd, J=2.4, 8.2Hz, 1H), 7.50 (d, J=8.2Hz, 1H), 6.68 (s, 2H), 3.86 (s, 2H), 3.13 (s, 4H), 1.02 (s, 6H) in CD <sub>3</sub> OD

TABLE 11:

No.	Chemical Structure	Salt	Properties m.p. (°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR (DMSO-d <sub>6</sub> )
51		fumarate	milky white cryst. 120-122°C acetone	m/z 176 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>13</sub> N <sub>3</sub>	8.56 (d, J=4.7Hz, 1H), 7.84 (t, J=7.0, 7.8Hz, 1H), 7.41 (d, J=7.8Hz 1H), 7.37 (t, J=4.7, 7.0Hz, 1H), 6.70 (s, 2H), 3.95 (s, 2H), 3.46 (t, J=5.7Hz, 4H), 2.01 (q, J=5.7Hz, 2H) in CD <sub>3</sub> OD
52		fumarate	colorless cryst. 185-186°C acetone	m/z 244 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>11</sub> Cl <sub>2</sub> N <sub>3</sub>	8.37 (d, J=2.1Hz, 1H), 8.33 (s, 1H), 8.10 (d, J=2.1Hz, 1H), 6.68 (s, 2H), 4.70 (s, 2H), 3.31 (m, 4H), 2.04 (m, 2H) in CD <sub>3</sub> OD
53		fumarate	colorless cryst. 152°C acetone	m/z 244 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>11</sub> Cl <sub>2</sub> N <sub>3</sub>	8.36 (d, J=2.1Hz, 1H), 8.06 (d, J=2.1Hz, 1H), 6.71 (s, 2H), 4.01 (t, J=11.5Hz, 2H), 3.80 (t, J=11.5Hz, 2H), 3.34 (s, 2H), 3.20 (s, 3H) in CD <sub>3</sub> OD
54		fumarate	colorless cryst. 157°C acetone	m/z 224 = (M+H) <sup>+</sup> C <sub>11</sub> H <sub>14</sub> ClN <sub>3</sub>	8.37 (d, J=2.5Hz, 1H), 7.81 (m, 1H), 7.51 (m, 1H), 6.70 (s, 2H), 3.83 (s, 2H), 3.69 (m, 1H), 3.45 (m, 2H), 2.11 (m, 1H), 1.68 (m, 1H), 1.34 (m, 3H) in CD <sub>3</sub> OD
55		fumarate	pale yellow cryst. 138-143°C acetone	m/z 224 = (M+H) <sup>+</sup> C <sub>11</sub> H <sub>14</sub> ClN <sub>3</sub>	8.34 (s, 1H), 8.03 (s, 1H), 7.81 (d, J=8.1Hz, 1H), 7.50 (d, J=8.1Hz, 1H), 6.37 (s, 2H), 3.67 (t, J=6.9Hz, 2H), 3.42 (m, 2H), 3.22 (m, 2H), 2.95 (t, J=6.9Hz, 2H), 1.89 (m, 2H)



TABLE 12:






No.	Chemical Structure	Salt	Properties m.p. (°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR (DMSO-d <sub>6</sub> )
56		fumarate (1.5 molecules)	colorless cryst. 124–125°C acetone	m/z 177 = (M+H) <sup>+</sup> C <sub>9</sub> H <sub>12</sub> N <sub>4</sub>	9.22 (s, 1H), 8.37 (s, 1H), 7.80 (s, 1H), 7.79 (s, 1H), 6.71 (s, 3H), 5.01 (s, 2H), 3.49 (t, J=5.5Hz, 2H), 3.43 (t, J=5.5Hz, 2H), 2.11 (t, J=5.5Hz, 2H) in CD <sub>3</sub> OD
57		fumarate (2 molecules)	colorless cryst. 156–157°C acetone	m/z 190 = (M+H) <sup>+</sup> C <sub>11</sub> H <sub>16</sub> N <sub>3</sub>	8.49 (s, 2H), 7.72 (d, J=7.8Hz, 1H), 7.32 (d, J=7.8Hz, 1H), 6.53 (s, 4H), 4.65 (s, 2H), 3.25 (m, 4H), 2.50 (s, 3H), 1.87 (m, 2H)
58		fumarate (2 molecules)	colorless cryst. 141–142°C acetone	m/z 176 = (M+H) <sup>+</sup> C <sub>10</sub> H <sub>13</sub> N <sub>3</sub>	8.62 (s, 1H), 8.58 (d, J=4.8Hz, 1H), 8.49 (s, 1H), 7.83 (d, J=7.7Hz, 1H), 7.46 (dd, J=4.8, 7.7Hz, 1H), 6.52 (s, 4H), 4.69 (s, 2H), 3.25 (m, 4H), 1.87 (m, 2H)
59		hydrochloride (2 molecules)	yellow cryst. 134–140°C acetonitrile	m/z 211 = (M+H) <sup>+</sup> C <sub>4</sub> H <sub>11</sub> ClN <sub>4</sub>	11.46 (br, 1H), 10.21 (br, 1H), 8.47 (s, 1H), 7.93 (d, J=8.2Hz, 1H), 7.57 (d, J=8.2Hz, 1H), 5.94 (br, 1H), 3.81 (s, 2H), 3.38 (m, 2H), 3.00 (m, 2H)
60		fumarate	colorless cryst. 156–158°C acetone	m/z 224 = (M+H) <sup>+</sup> C <sub>11</sub> H <sub>14</sub> ClN <sub>3</sub>	8.37 (d, J=2.5Hz, 1H), 7.81 (dd, J=2.5, 8.3Hz, 1H), 7.50 (d, J=8.3Hz, 1H), 6.68 (s, 2H), 4.04 (q, J=7.2Hz, 1H), 3.45 (t, J=5.7Hz, 4H), 1.98 (quintet, J=5.7Hz, 2H), 1.63 (d, J=7.2Hz, 3H) in CD <sub>3</sub> OD

TABLE 13:









No.	Chemical Structure	Salt	Properties m.p. (°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR(DMSO-d <sub>6</sub> )
61		fumarate	colorless cryst. 133-134°C acetone	m/z 216 = (M+H) <sup>+</sup> C <sub>8</sub> H <sub>10</sub> ClN <sub>3</sub> S	8.11 (s, 1H), 7.66 (s, 1H), 6.41 (s, 2H), 4.56 (s, 2H), 3.35 (m, 4H), 1.77 (m, 2H)
62		fumarate	colorless cryst. 144-146°C acetone	m/z 224 = (M+H) <sup>+</sup> C <sub>11</sub> H <sub>14</sub> ClN <sub>3</sub>	8.38 (d, J=2.1Hz, 1H), 7.85 (dd, J=2.1, 8.2Hz, 1H), 7.50 (d, J=8.2Hz, 1H), 6.38 (s, 2H), 3.75 (m, 4H), 3.59 (t, J=7.2Hz, 2H), 2.91 (t, J=7.2Hz, 2H), 2.09 (s, 3H)
63		hydrochloride (2 molecules)	colorless cryst. 158-162°C acetone	m/z 238 = (M+H) <sup>+</sup> C <sub>12</sub> H <sub>16</sub> ClN <sub>3</sub>	10.34 (1H, s), 8.36 (d, J=2.4Hz, 1H), 8.28 (1H, s), 7.81 (dd, J=2.4, 8.2Hz, 1H), 7.52 (d, J=8.2Hz, 1H), 3.74 (t, J=6.8Hz, 4H), 3.62 (s, 2H), 2.97 (t, J=6.8Hz, 2H), 2.09 (s, 3H), 1.31 (s, 3H)
64		hydrochloride (2 molecules)	colorless cryst. 213-220°C acetone	m/z 216 = (M+H) <sup>+</sup> C <sub>8</sub> H <sub>10</sub> ClN <sub>3</sub> S	10.06 (s, 2H), 7.70 (s, 1H), 4.07 (s, 2H), 3.32 (m, 4H), 1.82 (m, 2H)
65		fumarate	yellow cryst. 148-150°C acetone	m/z 202 = (M+H) <sup>+</sup> C <sub>7</sub> H <sub>8</sub> ClN <sub>3</sub> S	7.58 (s, 1H), 6.49 (s, 2H), 4.03 (s, 2H), 3.65 (s, 4H)

TABLE 14:

No.	Chemical Structure	Salt	Properties m.p. (°C) crystallized solvent	Mass Spectrum found molecular formula	<sup>1</sup> H-NMR (DMSO-d <sub>6</sub> )
66		fumarate	colorless cryst. 151-156°C acetone	m/z 177 = (M+H) <sup>+</sup> C <sub>9</sub> H <sub>12</sub> N <sub>4</sub>	9.13 (s, 1H), 8.85 (s, 2H), 6.43 (s, 2H), 3.90 (s, 2H), 3.33 (m, 4H), 1.82 (m, 2H)
67		fumarate	colorless cryst. 155-156°C acetone	m/z 163 = (M+H) <sup>+</sup> C <sub>8</sub> H <sub>10</sub> N <sub>4</sub>	9.12 (s, 1H), 8.80 (s, 2H), 6.46 (s, 2H), 3.89 (s, 2H), 3.71 (s, 4H)
68		fumarate	colorless cryst. 137-139°C acetone	m/z 190 = (M+H) <sup>+</sup> C <sub>11</sub> H <sub>15</sub> N <sub>3</sub>	10.42 (s, 2H), 8.40 (s, 1H), 8.35 (s, 1H), 7.63 (s, 1H), 6.47 (s, 2H), 3.78 (s, 1H), 3.33 (m, 4H), 2.29 (s, 3H), 1.81 (m, 2H)

The effect of the compounds (I) of the present invention was evaluated by the following biological experiments.

Biological Experiment 1:

5 Binding assays at  $\alpha 4\beta 2$  subtype of nicotinic acetylcholine receptors

The affinity of the compounds of the present invention to  $\alpha 4\beta 2$  subtype of nicotinic acetylcholine receptors was performed by the following method, which was modified method described by  
10 Pabreza L. A., Dhawan S. & Kellar K. J., *Mol. Pharm.*, 39, 9-12 (1990), and by Anderson D. J. & Arneric S. P., *Eur. J. Pharm.*, 253, 261-267 (1994).

(1) Preparation of rat brain membrane containing  $\alpha 4\beta 2$  subtype of nicotinic acetylcholine receptors

15 Fischer-344 strain male rats (body weight: 200-240 g; 9 weeks old) obtained from Charles River Japan were used. Rats were housed in the breeding cage controlled of the room temperature at  $23 \pm 1^\circ\text{C}$ , and the humidity of  $55 \pm 5\%$  for 1 to 4 weeks. Rats (3 to 4 rats per a cage) were housed with lights on  
20 for 12 hours daily (from 7:00 to 19:00), and allowed free access to food and water.

Preparation of rat brain membrane containing  $\alpha 4\beta 2$  subtype of nicotinic acetylcholine receptors was performed as follow. That is, rat brains were isolated just after sacrificed by  
25 decapitation, washed with ice-cooled saline solution and then frozen at  $-80^\circ\text{C}$  with liquid nitrogen and stored till using. After thawing the frozen brain, the brain was homogenized in 10 volumes of ice-cooled buffer solution (50 mM of Tris-HCl, 120 mM of NaCl, 5 mM of KCl, 1 mM of  $\text{MgCl}_2$ , 2mM of  $\text{CaCl}_2$ ; pH 7.4;  $4^\circ\text{C}$ ) using  
30 homogenizer (HG30, Hitachi Kohki Ltd.) for 30 seconds, and the homogenate were centrifuged under  $1,000 \times G$  for 10 minutes at  $4^\circ\text{C}$ . The resulting supernatant was separated and the pellet was

homogenized again with half volume of aforementioned prior buffer solution and centrifuged under the same conditions. Combined supernatant was further centrifuged under 40,000 x G for 20 minutes at 4°C. The pellet was suspended in buffer solution and used for binding assays at receptors.

(2) Experiments of  $\alpha 4\beta 2$  subtype of nicotinic acetylcholine receptors binding

Suspensions of membrane pellets containing 400-600  $\mu\text{g}$  of protein were added to test tubes containing test compounds and [ $^3\text{H}$ ]-cytisine (2 nM) in a final volume of 200  $\mu\text{l}$  and incubated for 75 minutes in ice-cooled bath. The samples were isolated by vacuum filtration onto Whatman GF/B filters, which were prerinsed with 0.5% polyethylenimine just prior to sample filtration, using Brandel multi manifold cell harvester. The filters were rapidly washed with buffer solution (3 x 1 ml). The filters were counted in 3 ml of clearsol I (Nacalai Tesque Inc.). The determination of nonspecific binding was incubated in the presence of 10  $\mu\text{M}$  (-)-nicotine.

The analyses of the experimental results were conducted by using the Accufit Competition Program (Beckman Ltd.).

Biological Experiment 2:

Binding assays at  $\alpha 1\beta 1\gamma \delta$  subtype of nicotinic acetylcholine receptors

The affinity of the compounds of the present invention to  $\alpha 1\beta 1\gamma \delta$  subtype of nicotinic acetylcholine receptors was measured by the following method, which was modified method described by Garcha H. S., Thomas P., Spivak C. E., Wonnacott S. & Stolerman I. P., *Psychropharmacology*, 110, 347-354 (1993).

(1) Preparation of rat skeletal muscles containing  $\alpha 1\beta 1\gamma \delta$  subtype of nicotinic acetylcholine receptors

The substantially same animals described in the Biological Experiment 1 were used.

The isolation of  $\alpha 1\beta 1\gamma\delta$  subtype of nicotinic acetylcholine receptors was performed as follow. That is, rat posterior skeletal muscles were isolated just after sacrificed by decapitation, washed with ice-cooled saline solution and then frozen at  $-80^{\circ}\text{C}$  with liquid nitrogen and stored till using. After thawing the frozen muscles, tissue was homogenized (40% w/v) with buffer solution [2.5 mM of sodium phosphate buffer (pH:7.2), 90 mM of NaCl, 2 mM of KCl, 1 mM of EDTA, 2 mM of benzamidine, 0.1 mM of benzethonium chloride, 0.1 mM of PMSF, 0.01% of sodium azide] in Waring blender (Waring blender 34BL97; WARING PRODUCTS DIVISION DYNAMICS CORPORATION OF AMERICA) for 60 seconds. The homogenate were centrifuged under  $20,000 \times G$  for 60 minutes at  $4^{\circ}\text{C}$ . The supernatant was separated and the resulting pellet was added to the same buffer (1.5 ml/g wet weight), and homogenized under the same conditions. Triton X100 (2% w/v) was added and the mixture was stirred for 3 hours at  $4^{\circ}\text{C}$ . The centrifugation at  $100,000 \times G$  for 60 minutes at  $4^{\circ}\text{C}$  yielded the rat muscle extract as supernatant. This was stored at  $4^{\circ}\text{C}$  for up to 4 weeks, and used for binding assays at receptors.

(2) Experiments of  $\alpha 1\beta 1\gamma\delta$  subtype of nicotinic acetylcholine receptors binding

Receptors binding experiments were performed as follow. That is, the extract of rat muscle containing 600-900  $\mu\text{g}$  of protein was added to test tubes containing test compounds and incubated for 15 minutes at  $37^{\circ}\text{C}$ . Then, to this mixture was added 1 nM of [ $^3\text{H}$ ]- $\alpha$ -bungarotoxin ( $\alpha$ -Bgt) and further incubated for 2 hours. The samples were isolated by vacuum filtration onto Whatman GF/B filters, which were prerinsed with 0.5% polyethylenimine just prior to sample filtration, using Brandel

multi manifold cell harvester. The filters were rapidly rinsed with washing solution (10 mM of  $\text{KH}_2\text{PO}_4$ , 150 mM of NaCl, pH 7.2, room temperature) (5 x 1 ml). The filters were counted in 3 ml of clearsol I (Nacalai Tesque Inc.). The determination of  
5 nonspecific binding was incubated in the presence of 1  $\mu\text{M}$   $\alpha$ -Bgt. The solutions containing  $\alpha$ -Bgt (labeled/non-labeled) were prepared by using buffer solution containing 0.25% of BSA. In the receptor binding experiments, said buffer solution was added for adjusting the final concentration of BSA to be 0.05%.

10 The analyses of the experimental results were conducted by the same way as described in the Biological Experiment 1.

Table 15 shows the results of receptor binding studies of the compounds of the present invention and (-)-nicotine as  
15 reference compound.

TABLE 15:

Compound No.	Affinities for receptors Ki	
	$\alpha 4\beta 2$	$\alpha 1\beta 1\gamma\delta$ *1
2	13 nM	(34%, 6%)
3	45 nM	(34%, 5%)
4	67 nM	(46%, 16%)
7	86 nM	(80%, 51%)
8	29 nM	395 $\mu$ M
9	7.7 nM	(43%, 16%)
10	11 nM	(40%, 17%)
11	115 nM	(74%, 53%)
12	268 nM	(79%, 42%)
15	950 nM	n.d.
16	392 nM	(63%, 30%)
18	86 nM	(62%, 18%)
19	144 nM	(69%, 29%)
22	429 nM	(23%, -4%)
25	338 nM	(41%, 7%)
27	2 nM	45 $\mu$ M
32	580 nM	(69%, 53%)
33	365 nM	n.d.
36	124 nM	(81%, 34%)
43	167 nM	(71%, 28%)
48	82 nM	257 $\mu$ M
49	211 nM	773 $\mu$ M
52	1.2 nM	23 $\mu$ M
53	10 nM	83 $\mu$ M
54	108 nM	1739 $\mu$ M
57	12 nM	86 $\mu$ M
58	6.9 nM	32 $\mu$ M
62	70 nM	639 $\mu$ M
64	8.1 nM	23 $\mu$ M
65	53 nM	524 $\mu$ M
66	90 nM	841 $\mu$ M
68	203 nM	231 $\mu$ M
Nicotine	1.6 nM	182 $\mu$ M

\*1: Values indicated in a parenthesis show control % of [ $^3$ H]- $\alpha$ -Bgt binding at 100  $\mu$ M and 1,000  $\mu$ M of test compounds.

5 n.d.: not determined.



Biological Experiment 3:Agonist activities at human  $\alpha 4\beta 2$  subtype of nicotinic acetylcholine receptors

The agonist activities of the compounds of the present invention at human  $\alpha 4\beta 2$  subtype of nicotinic acetylcholine receptors was evaluated by the following method, which was modified method described by Papke R. L., Thinschmidt J. S., Moulton B. A., Meyer E. M. & Poirier A., *Br. J. Pharmacol.*, 120, 429-438 (1997).

10 (1) Preparation of cRNA of human  $\alpha 4\beta 2$  subtype of nicotinic acetylcholine receptors

The cloning of human nicotinic acetylcholine receptor (hnACh-R)  $\alpha 4$  cDNA and hnACh-R  $\beta 2$  cDNA were performed, in accordance with the conventional manners, by synthesizing the each DNA primers corresponding to the sequences of hnACh-R  $\alpha 4$  cDNA and hnACh-R  $\beta 2$  cDNA [Monteggia L. M. et al., *Gene*, 155, 189-193 (1995); and Anand R., & Lindstrom J., *Nucl. Acids Res.*, 18, 4272 (1990)], and obtained hnACh-R  $\alpha 4$  cDNA and hnACh-R  $\beta 2$  cDNA by polymerase chain reaction (PCR), respectively. The obtained hnACh-R  $\alpha 4$  cDNA and hnACh-R  $\beta 2$  cDNA were inserted to the cRNA expression vector (pSP64 polyA) having SP6 RNA promoter to construct hnACh-R  $\alpha 4$ /pSP64 polyA and hnACh-R  $\beta 2$ /pSP64 polyA, respectively. After cutting from expression vector by restriction enzyme (EcoRI), transcription was performed by affecting SP6 RNA polymerase in the presence of cap analogues to obtain hnACh-R  $\alpha 4$  cRNA and hnACh-R  $\beta 2$  cRNA, respectively.

20 (2) Expression of human  $\alpha 4\beta 2$  subtype nicotinic acetylcholine receptors in *Xenopus* oocytes

30 Oocytes were purchased from Kitanihonseibutsukyohzai Co., Ltd., which were already enucleated from *Xenopus laevis*, and used in this experiment.

The oocytes were treated with collagenase (Sigma type I; 1 mg/ml) in calcium-free modified BIRTH's solution (88 mM of NaCl, 1 mM of KCl, 2.4 mM of NaHCO<sub>3</sub>, 0.82 mM of MgSO<sub>4</sub>, 15 mM of HEPES, pH 7.6) under gently stirring at room temperature for 90 minutes, and washed out the enzyme from the tissue. Then, oocytes were separated from ovarian follicle by tweezers, and isolated oocytes were placed in antibiotics containing modified BIRTH's solution (88 mM of NaCl, 1 mM of KCl, 2.4 mM of NaHCO<sub>3</sub>, 0.82 mM of MgSO<sub>4</sub>, 15 mM of HEPES, pH 7.6, and 0.1 v/v% of mixture solution containing of penicillin and streptomycin for incubation; Sigma Co.). Thus treated oocytes were injected with 50 nl of adjusted cRNAs (1.0 mg/ml), that is, each 50 ng of hnACh-R  $\alpha 4$  cRNA and hnACh-R  $\beta 2$  cRNA per 1 oocyte by using automatic injector (NANOJECT; DRUMMOND SCIENTIFIC CO.), and further incubated for 4-14 days at 19°C. In oocytes, heterogeneous quintuple  $[(\alpha 4)_2(\beta 2)_3]$  was composed by translation of injected cRNAs, and ion channel receptors were constructed on cell membrane.

(3) Agonist activities at human  $\alpha 4\beta 2$  subtype of nicotinic acetylcholine receptors

The recordings of responses at human  $\alpha 4\beta 2$  subtype of nicotinic acetylcholine receptors by means of membrane potential holding method were performed as follow. That is, oocytes were placed in recording chamber with a total volume of 50  $\mu$ l and were perfused with Ringer's solution (115 mM of NaCl, 2.5 mM of KCl, 1.8 mM of CaCl<sub>2</sub>, 10 mM of HEPES, pH 7.3) containing atropine (1  $\mu$ M) under flow rate of 1 ml/min. The membrane electric potentials were held at -50 mV by mean of the two electric membranes potential holding method (CEZ-1250; Nihon Kohden Co.). Test compounds were added to the perfusion solution, and recorded the peak strength of induced inward current. In order to normalize the responses of test compounds, the responses with acetylcholine

(ACh) were recorded before and after application of the test compounds. Generally in the oocytes just after isolated, the response of intrinsic muscarinic acetylcholine receptors, which is inward current caused by activation of calcium dependence chloride ion channels with increase of the intracellular calcium concentration by stimulation of receptors, is observed. However, the complete disappearances of these responses were confirmed when treated with collagenase or added 1  $\mu$ M of atropine. Furthermore, the oocytes without injection of cRNAs showed no responses by ACh after treatment with collagenase. Therefore, the responses observed in oocytes with injection of hnACh-R  $\alpha 4$  cRNA and hnACh-R  $\beta 2$  cRNA, i.e., the inward current induced by the intracellular influx of sodium ion according to the stimulation of receptors, would be the freshly observed responses of human  $\alpha 4\beta 2$  subtype nicotinic acetylcholine receptors.

Table 16 shows the results of the agonist activity test of the compounds in the present invention and (-)-nicotine as reference compound.

TABLE 16:

Compound No.	Agonist activity (ED50) <sup>*1</sup>
2	3.4 $\mu$ M
3	43.8 $\mu$ M
22	(13.2%)
27	(18.0%)
45	(12.0%)
57	(9.1%)
58	(27.9%)
62	(9.6%)
nicotine	11.4 $\mu$ M

<sup>\*1</sup>: These data are calculated in comparison with the reaction with 10  $\mu$ M of acetylcholine (100%). Values indicated in a parenthesis show control % by response at 100  $\mu$ M of the test compounds.

The following are Formulation Examples of the compounds (I) or pharmaceutically acceptable salt thereof according to the present invention

Formulation Example 1 (Tablets):

Compound 2 (Fumarate)	25 g
Lactose	130 g
Crystalline cellulose	20 g
Corn starch	20 g
3% aqueous solution of hydroxypropylmethyl-cellulose	100 ml
Magnesium stearate	2 g

Fumarate of Compound 2, lactose, crystalline cellulose and corn starch were screened through a 60-mesh sieve, homogenized and charged into a kneader. A 3% aqueous solution of hydroxypropylmethylcellulose was added to the homogeneous mixture and the mixture was further kneaded. The product was granulated by a 16-mesh sieve, dried in air at 50°C, and again granulated by

a 16-mesh sieve. Magnesium stearate was added to the granule and mixed again. The mixture was tabletted to produce tablets weighing 200 mg each and having an 8 mm diameter.

5 Formulation Example 2 (Capsules):

Compound 3 (Fumarate)	25.0 g
Lactose	125.0 g
Corn starch	48.5 g
Magnesium stearate	1.5 g

10 The above components were finely pulverized and thoroughly mixed to produce a homogeneous mixture. The mixture was filled in gelatin capsules, 200 mg per capsule, to obtain capsules.

Formulation Example 3 (Injection):

15 The fumarate of Compound 58 was filled in an amount of 250 mg in a vial and mixed in situ with approximately 4-5 ml of injectable distilled water to make an injectable solution.

INDUSTRIAL APPLICABILITY


20 As described above, the compounds of the present invention possess high affinity for  $\alpha 4\beta 2$  nicotinic acetylcholine receptor of central nervous systems and activate said  $\alpha 4\beta 2$  nicotinic acetylcholine receptor as agonists or modulators. Therefore, the compounds of the present invention are useful for preventing or  
25 treating various kinds of diseases, which may be prevented or cured by activating nicotinic acetylcholine receptors.

Especially, the activators for  $\alpha 4\beta 2$  nicotinic acetylcholine receptors of the present invention are useful for preventing or treating various diseases such as dementia, senile  
30 dementia, presenile dementia, Alzheimer's disease, Parkinson's disease, cerebrovascular dementia, AIDS-related dementia, dementia in Down's syndrome, Tourette's syndrome, neurosis during

the chronic cerebral infarction stage, cerebral dysfunction caused by cerebral injury, anxiety, schizophrenia, depression, Huntington's disease, pain and so on.

## CLAIMS

1. Cyclic amidine compounds represented by the following formula (I):



wherein:

A¹ and A² are hydrogen atom, optionally substituted alkyl group; optionally substituted aryl group; or optionally substituted heterocyclic group; and

X is -C(R¹,R²)-C(R³,R⁴)-, -C(R⁵)=C(R⁶)-, -C(R⁷,R⁸)-C(R⁹,R¹⁰)-C(R¹¹,R¹²)-, or -C(R¹³,R¹⁴)-C(R¹⁵,R¹⁶)-NH- (wherein, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵ and R¹⁶ are hydrogen atom; halogen atom; optionally substituted alkyl group; optionally substituted aryl group; or optionally substituted heterocyclic group; or pharmaceutically acceptable salts thereof.

2. The following compounds represented by the formula (I) of claim 1;

- 2-(6-chloro-3-pyridyl)-2-imidazoline;
- 2-(6-chloro-3-pyridyl)-1,4,5,6-tetrahydropyrimidine;
- 2-(6-chloro-3-pyridyl)-1-methyl-2-imidazoline;
- 2-(6-chloro-3-pyridyl)-1-methyl-1,4,5,6-tetrahydropyrimidine;
- 1-(6-chloro-3-pyridyl)methylimidazole;
- 2-(6-chloro-3-pyridyl)imidazole;
- 2-(6-chloro-3-pyridyl)methyl-2-imidazoline;
- 2-(6-chloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- 2-(6-chloro-3-pyridyl)methyl-1-methyl-2-imidazoline;
- 2-(6-chloro-3-pyridyl)methyl-1-methyl-1,4,5,6-

- tetrahydropyrimidine;
- 1-(6-chloro-3-pyridyl)methyl-2-methyl-2-imidazoline;
- 1-(6-chloro-3-pyridyl)methyl-4,4-dimethyl-2-imidazoline;
- 2-(tetrahydrofuran-3-yl)-1,4,5,6-tetrahydropyrimidine;
- 5 2-(tetrahydrofuran-3-yl)-2-imidazoline;
- 2-(tetrahydrofuran-3-yl)methyl-1,4,5,6-tetrahydropyrimidine;
- 2-(5-bromo-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- 2-(5-bromo-3-pyridyl)methyl-2-imidazoline;
- 2-(3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- 10 2-(3-pyridyl)methyl-2-imidazoline;
- 2-(3-aminophenyl)-1,4,5,6-tetrahydropyrimidine;
- 2-(3-quinolyl)methyl-1,4,5,6-tetrahydropyrimidine;
- 2-(2-chloro-5-thiazolyl)-1,4,5,6-tetrahydropyrimidine;
- 2-(3-quinolyl)methyl-2-imidazoline;
- 15 2-(2-chloro-5-thiazolyl)-2-imidazoline;
- 2-(3-quinolyl)-1,4,5,6-tetrahydropyrimidine;
- 2-(3-furanyl)methyl-2-imidazoline;
- 1-(6-chloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- 2-(3,5-dimethyl-4-isoxazolyl)methyl-1,4,5,6-tetrahydro-
- 20 pyrimidine;
- 2-(3,5-dimethyl-4-isoxazolyl)methyl-2-imidazoline;
- 2-(3-thienyl)methyl-1,4,5,6-tetrahydropyrimidine;
- 2-(3-thienyl)methyl-2-imidazoline;
- 2-methyl-5-(3-pyridyl)-2-imidazoline;
- 25 5-(3-pyridyl)-2-imidazoline;
- 1,2-bis[(6-chloro-3-pyridyl)methyl]-1,4,5,6-tetrahydro-
- pyrimidine;
- 1-(6-chloro-3-pyridyl)methyl-2-(3-pyridyl)-2-imidazoline;
- 2-(5,6-dichloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;
- 30 2-(6-chloro-3-pyridyl)methyl-5-phenyl-1,4,5,6-tetrahydro-
- pyrimidine;
- 2-(4-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;



- 2-(2-chloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;  
2-(2,6-dichloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;  
2-[2-(6-chloro-3-pyridyl)ethyl]-1,4,5,6-tetrahydropyrimidine;  
2-[2-(6-chloro-3-pyridyl)ethyl]-2-imidazoline;  
5 2-(6-methyl-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;  
1,2-bis[(6-chloro-3-pyridyl)methyl]-2-imidazoline;  
2-(6-methyl-3-pyridyl)methyl-2-imidazoline;  
2-(6-ethoxy-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;  
2-(6-ethoxy-3-pyridyl)methyl-2-imidazoline;  
10 2-(6-fluoro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;  
2-(5,6-dichloro-3-pyridyl)methyl-2-imidazoline;  
2-(6-chloro-3-pyridyl)methyl-5,5-dimethyl-1,4,5,6-tetrahydro-  
pyrimidine;  
2-(2-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;  
15 1-(5,6-dichloro-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;  
2-(5,6-dichloro-3-pyridyl)methyl-1-methyl-2-imidazoline;  
2-(6-chloro-3-pyridyl)methyl-4-methyl-1,4,5,6-tetrahydro-  
pyrimidine;  
1-[2-(6-chloro-3-pyridyl)ethyl]-1,4,5,6-tetrahydropyrimidine;  
20 1-(3-pyridazinyl)methyl-1,4,5,6-tetrahydropyrimidine;  
1-(6-methyl-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;  
1-(3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine;  
3-(6-chloro-3-pyridyl)methyl-1,4,5,6-tetrahydro-1,2,4-triazine;  
2-[1-(6-chloro-3-pyridyl)ethyl]-1,4,5,6-tetrahydropyrimidine;  
25 1-(2-chloro-5-thiazolyl)methyl-1,4,5,6-tetrahydropyrimidine;  
1-[2-(6-chloro-3-pyridyl)ethyl]-2-methyl-2-imidazoline;  
1-[2-(6-chloro-3-pyridyl)ethyl]-4,4-dimethyl-2-imidazoline;  
2-(2-chloro-5-thiazolyl)methyl-1,4,5,6-tetrahydropyrimidine;  
2-(2-chloro-5-thiazolyl)methyl-2-imidazoline;  
30 2-(5-pyrimidyl)methyl-1,4,5,6-tetrahydropyrimidine;  
2-(5-pyrimidyl)methyl-2-imidazoline;  
2-(5-methyl-3-pyridyl)methyl-1,4,5,6-tetrahydropyrimidine.

and pharmaceutically acceptable salt thereof.

3. Activators for  $\alpha 4\beta 2$  nicotinic acetylcholine receptors containing the compound or pharmaceutically acceptable salt thereof claimed in claim 1 or 2, as active ingredient.

4. The activators for  $\alpha 4\beta 2$  nicotinic acetylcholine receptors according to claim 3, wherein said activators are agonists or modulators at  $\alpha 4\beta 2$  nicotinic acetylcholine receptors.

5. A medicament for preventing or treating cerebral circulation diseases comprising the activator for  $\alpha 4\beta 2$  nicotinic acetylcholine receptors claimed in claim 3 or 4.

6. A medicament for preventing or treating neurodegenerative disease, dementia, motor ataxia, and neuropathy and mental disease comprising the activator for  $\alpha 4\beta 2$  nicotinic acetylcholine receptors claimed in claim 3 or 4.

7. The medicament according to claim 6, wherein said neurodegenerative disease is Alzheimer's disease or Parkinson's disease, said dementia is cerebrovascular dementia, said motor ataxia is Tourette's syndrome, and said neuropathy and mental disease is neurosis during the chronic cerebral infarction stage, anxiety or schizophrenia.

8. A medicament for improving the cerebral metabolism, neurotransmission functional disorder and memory disorder, for protecting brain, or having analgesic effect, which comprises the activator for  $\alpha 4\beta 2$  nicotinic acetylcholine receptors claimed in claim 3 or 4.

9. A medicament for preventing or treating inflammatory intestinal diseases comprising the activator for  $\alpha 4\beta 2$  nicotinic acetylcholine receptors claimed in claim 3 or 4.

5 10. The use of the compounds claimed in claim 1 or 2 as the activators for  $\alpha 4\beta 2$  nicotinic acetylcholine receptors.

11. The method of preventing or treating cerebral circulation diseases which comprises administering activators for  $\alpha 4\beta 2$   
10 nicotinic acetylcholine receptors claimed in claim 3 or 4.

12. The method of preventing or treating neurodegenerative diseases, dementia, motor ataxia, and neuropathy and mental disease which comprises administering activators for  $\alpha 4\beta 2$   
15 nicotinic acetylcholine receptors claimed in claim 3 or 4.

13. The method according to claim 12, wherein said neurodegenerative disease is Alzheimer's disease or Parkinson's disease, said dementia is cerebrovascular dementia, said motor  
20 ataxia is Tourette's syndrome, and said neuropathy and mental disease is neurosis during the chronic cerebral infarction stage, anxiety or schizophrenia.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



10/009433

(43) International Publication Date  
1 November 2001 (01.11.2001)

PCT

(10) International Publication Number  
WO 01/081334 A3

(51) International Patent Classification<sup>7</sup>: C07D 401/04,  
A61K 31/395, A61P 43/00, C07D 401/06, 405/04, 405/06,  
239/06, 417/04, 413/06, 409/06, 401/14, 417/06, 237/26

AKABANE, Minako [JP/JP]; 1-37-201, Sojiji 1-chome,  
Ibaraki-shi, Osaka 567-0801 (JP). TANI, Yoshihiro  
[JP/JP]; 9-38, Hoshimi-cho, Ibaraki-shi, Osaka 567-0843  
(JP).

(21) International Application Number: PCT/JP01/03378

(22) International Filing Date: 20 April 2001 (20.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
2000-120976 21 April 2000 (21.04.2000) JP

(74) Agent: KUSAMA, Osamu; KUSAMA PATENT  
OFFICE, 7F Iwata Bldg., 5-12, Iidabashi 4-chome, Chiyo-  
oda-ku, Tokyo 102-0072 (JP).

(81) Designated States (*national*): AU, CA, CN, KR, US.

(84) Designated States (*regional*): European patent (AT, BE,  
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,  
NL, PT, SE, TR).

(71) Applicant (*for all designated States except US*): SUN-  
TORY LIMITED [JP/JP]; 1-40, Dojimahama 2-chome,  
Kita-ku, Osaka-shi, Osaka 530-8203 (JP).

Published:  
— with international search report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): IMOTO, Masahiro  
[JP/JP]; 15-501, Naka-machi 3-chome, Tatebayashi-shi,  
Gunma 374-0029 (JP). IWANAMI, Tatsuya [JP/JP];  
16-16, Dai-cho, Ashikaga-shi, Tochigi 326-0056 (JP).

(88) Date of publication of the international search report:  
8 August 2002


*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

RECEIVED

SEP 06 2002

TECH CENTER 1600/2900

(54) Title: CYCLIC AMIDINE COMPOUNDS AND THEIR USE AS ALPHA4BETA2 NICOTINIC ACETYLCHOLINE RE-  
CEPTOR LIGANDS



(I)

(57) Abstract: There is provided cyclic amidine compounds of the following formula (I) wherein: A<sup>1</sup> and A<sup>2</sup> are hydrogen atom, optionally substituted alkyl group; optionally substituted aryl group; or optionally substituted heterocyclic group; and X is -C(R<sup>1</sup>,R<sup>2</sup>)-C(R<sup>3</sup>,R<sup>4</sup>)-, -C(R<sup>5</sup>)=C(R<sup>6</sup>)-, -C(R<sup>7</sup>,R<sup>8</sup>)-C(R<sup>9</sup>,R<sup>10</sup>)-C(R<sup>11</sup>,R<sup>12</sup>)-, or -C(R<sup>13</sup>,R<sup>14</sup>)-C(R<sup>15</sup>,R<sup>16</sup>)-NH- (wherein, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup>, R<sup>14</sup>, R<sup>15</sup> and R<sup>16</sup> are hydrogen atom; halogen atom; optionally substituted alkyl group; optionally substituted aryl group; or optionally substituted heterocyclic group; or pharmaceutically acceptable salts thereof. These

compounds have good affinity for α4β2 nicotinic acetylcholine receptors and activate the same to thereby exert a preventive or therapeutic effect on cerebral dysfunction.

WO 01/081334 A3

## INTERNATIONAL SEARCH REPORT

 Int: al Application No  
 PCT/JP 01/03378

## A. CLASSIFICATION OF SUBJECT MATTER

 IPC 7 C07D401/04 A61K31/395 A61P43/00 C07D401/06 C07D405/04  
 C07D405/06 C07D239/06 C07D417/04 C07D413/06 C07D409/06  
 C07D401/14 C07D417/06 C07D237/26

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, WPI Data, EPO-Internal

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LATLI, BACHIR ET AL: "Novel and Potent 6-Chloro-3-pyridinyl Ligands for the.alpha.4.beta.2 Neuronal Nicotinic Acetylcholine Receptor" J. MED. CHEM. (1999), 42(12), 2227-2234 , XP000941733 cited in the application compound 12 table 2  --- -/---	1-13

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

9 January 2002

Date of mailing of the international search report

25/01/2002

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel: (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

De Jong, B

## INTERNATIONAL SEARCH REPORT

Internal Application No  
PCT/JP 01/03378

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; AKEMASA, HIROYUKI: "Preparation of pyridine derivatives as intermediates for insecticides and their intermediates" retrieved from STN Database accession no. 120:298481 XP002185104 abstract & JP 05 339267 A (KOEI CHEMICAL CO, JAPAN) 21 December 1993 (1993-12-21) ----	1,2
A	EP 0 679 397 A (BAYER AG) 2 November 1995 (1995-11-02) cited in the application claim 1 ----	1-13
A	WO 00 10997 A (ORTHO MCNEIL PHARM INC) 2 March 2000 (2000-03-02) page 30, line 10 - line 13 -----	1,3-13

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1,3-13

Present claims 1,3-13 relate to an extremely large number of possible compounds, their use and pharmaceuticals containing them. Claim 1 even covers well known compounds such as imidazole, 2-imidazoline and 1,4,5,6-tetrahydropyrimidine. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search is only complete for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds according to claim 2, their use and pharmaceuticals containing them.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/JP 01/03378

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
JP 5339267	A	21-12-1993	JP	3179578 B2	25-06-2001
EP 0679397	A	02-11-1995	DE	4414569 A1	02-11-1995
			CA	2147930 A1	28-10-1995
			EP	0679397 A2	02-11-1995
			JP	7300415 A	14-11-1995
			US	5547965 A	20-08-1996
WO 0010997	A	02-03-2000	AU	5574799 A	14-03-2000
			EP	1107965 A1	20-06-2001
			WO	0010997 A1	02-03-2000



## PCT

## INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference <b>SN-45</b>	<b>FOR FURTHER ACTION</b> see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. <b>PCT/JP 01/03378</b>	International filing date (day/month/year) <b>20/04/2001</b>	(Earliest) Priority Date (day/month/year) <b>21/04/2000</b>
Applicant <b>SUNTORY LIMITED</b>		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 5 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

## 1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.

☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing:

☐ contained in the international application in written form.

☐ filed together with the international application in computer readable form.

☐ furnished subsequently to this Authority in written form.

☐ furnished subsequently to this Authority in computer readable form.

☐ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.

☐ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☒ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,

☐ the text is approved as submitted by the applicant.

☒ the text has been established by this Authority to read as follows:

**CYCLIC AMIDINE COMPOUNDS AND THEIR USE AS ALPHA4BETA2 NICOTINIC ACETYLCHOLINE RECEPTOR LIGANDS**

5. With regard to the **abstract**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.

☐ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

☒ None of the figures.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1,3-13

Present claims 1,3-13 relate to an extremely large number of possible compounds, their use and pharmaceuticals containing them. Claim 1 even covers well known compounds such as imidazole, 2-imidazoline and 1,4,5,6-tetrahydropyrimidine. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search is only complete for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds according to claim 2, their use and pharmaceuticals containing them.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP 01/03378

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/04 A61K31/395 A61P43/00 C07D401/06 C07D405/04  
C07D405/06 C07D239/06 C07D417/04 C07D413/06 C07D409/06  
C07D401/14 C07D417/06 C07D237/26

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, WPI Data, EPO-Internal

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LATLI, BACHIR ET AL: "Novel and Potent 6-Chloro-3-pyridinyl Ligands for the.alpha.4.beta.2 Neuronal Nicotinic Acetylcholine Receptor" J. MED. CHEM. (1999), 42(12), 2227-2234 , XP000941733 cited in the application compound 12 table 2  --- -/--	1-13



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

\* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&amp;\* document member of the same patent family

Date of the actual completion of the international search

9 January 2002

Date of mailing of the international search report

25/01/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

De Jong, B

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/JP 01/03378

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE CA 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; AKEMASA, HIROYUKI: "Preparation of pyridine derivatives as intermediates for insecticides and their intermediates" retrieved from STN Database accession no. 120:298481 XP002185104 abstract & JP 05 339267 A (KOEI CHEMICAL CO, JAPAN) 21 December 1993 (1993-12-21) ---	1,2
A	EP 0 679 397 A (BAYER AG) 2 November 1995 (1995-11-02) cited in the application claim 1 ---	1-13
A	WO 00 10997 A (ORTHO MCNEIL PHARM INC) 2 March 2000 (2000-03-02) page 30, line 10 - line 13 -----	1,3-13

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 01/03378

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
JP 5339267	A	21-12-1993	JP 3179578 B2	25-06-2001
EP 0679397	A	02-11-1995	DE 4414569 A1	02-11-1995
			CA 2147930 A1	28-10-1995
			EP 0679397 A2	02-11-1995
			JP 7300415 A	14-11-1995
			US 5547965 A	20-08-1996
WO 0010997	A	02-03-2000	AU 5574799 A	14-03-2000
			EP 1107965 A1	20-06-2001
			WO 0010997 A1	02-03-2000

## PCT

NOTICE INFORMING THE APPLICANT OF THE  
COMMUNICATION OF THE INTERNATIONAL  
APPLICATION TO THE DESIGNATED OFFICES

(PCT Rule 47.1(c), first sentence)

From the INTERNATIONAL BUREAU

To:

KUSAMA, Osamu  
KUSAMA PATENT OFFICE  
7F Iwata Bldg.  
5-12, Iidabashi 4-chome  
Chiyoda-ku, Tokyo 102-0072  
JAPON

Date of mailing (day/month/year) 01 November 2001 (01.11.01)		IMPORTANT NOTICE	
Applicant's or agent's file reference SN-45			
International application No. PCT/JP01/03378	International filing date (day/month/year) 20 April 2001 (20.04.01)	Priority date (day/month/year) 21 April 2000 (21.04.00)	
Applicant SUNTORY LIMITED et al			

1. Notice is hereby given that the International Bureau has **communicated**, as provided in Article 20, the international application to the following designated Offices on the date indicated above as the date of mailing of this notice:  
**KR,US**

In accordance with Rule 47.1(c), third sentence, those Offices will accept the present notice as conclusive evidence that the communication of the international application has duly taken place on the date of mailing indicated above and no copy of the international application is required to be furnished by the applicant to the designated Office(s).

2. The following designated Offices have waived the requirement for such a communication at this time:  
**AU,CA,CN,EP**

The communication will be made to those Offices only upon their request. Furthermore, those Offices do not require the applicant to furnish a copy of the international application (Rule 49.1(a-bis)).

3. Enclosed with this notice is a copy of the international application as published by the International Bureau on  
01 November 2001 (01.11.01) under No. WO 01/81334

**REMINDER REGARDING CHAPTER II (Article 31(2)(a) and Rule 54.2)**

If the applicant wishes to postpone entry into the national phase until 30 months (or later in some Offices) from the priority date, a **demand for international preliminary examination** must be filed with the competent International Preliminary Examining Authority before the expiration of 19 months from the priority date.

It is the applicant's sole responsibility to monitor the 19-month time limit.

Note that only an applicant who is a national or resident of a PCT Contracting State which is bound by Chapter II has the right to file a demand for international preliminary examination (at present, all PCT Contracting States are bound by Chapter II).

**REMINDER REGARDING ENTRY INTO THE NATIONAL PHASE (Article 22 or 39(1))**

If the applicant wishes to proceed with the international application in the **national phase**, he must, within 20 months or 30 months, or later in some Offices, perform the acts referred to therein before each designated or elected Office.

For further important information on the time limits and acts to be performed for entering the national phase, see the Annex to Form PCT/IB/301 (Notification of Receipt of Record Copy) and the PCT Applicant's Guide, Volume II.

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland	Authorized officer  J. Zahra
Facsimile No. (41-22) 740.14.35	Telephone No. (41-22) 338.91.11